

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
31 January 2002 (31.01.2002)

PCT

(10) International Publication Number
WO 02/08263 A2

(51) International Patent Classification⁷: **C07K 14/005**

(21) International Application Number: **PCT/GB01/03252**

(22) International Filing Date: **19 July 2001 (19.07.2001)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
0017720.4 19 July 2000 (19.07.2000) **GB**

(71) Applicant (for all designated States except US): **GOT-A-GENE AB** [SE/SE]; Stena Center 1B, S-412 92 Göteborg (SE).

(71) Applicant (for GB only): **GARDNER, Rebecca** [GB/GB]; Frank B. Dehn & Co., 179 Queen Victoria Street, London EC4V 4EL (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **LINDHOLM, Leif** [SE/SE]; Got-A-Gene AB, Stena Center 1B, S-412 92 Göteborg (SE). **NORD, Anna, Karin** [SE/SE]; Af-fibody AB, Björnåsvägen 21, S-113 47 Stockholm (SE). **BOULANGER, Pierre, Alain** [FR/FR]; Faculté de Médecine RTH Laennec, Laboratoire de Virologie et Pathogénèse Virale CNRS UMR 5537, 7, rue Guillaume Paradin, F-69372 Lyon Cedex 8 (FR).

(74) Agents: **GARDNER, Rebecca** et al.; Frank B. Dehn & Co., 179 Queen Victoria Street, London EC4V 4EL (GB).

(81) Designated States (national): AE, AG, AL, AM, AT, AT (utility model), AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, CZ (utility model), DE, DE (utility model), DK, DK (utility model), DM, DZ, EC, EE, EE (utility model), ES, FI, FI (utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ,

[Continued on next page]

(54) Title: **MODIFIED VIRUS**

A. WT Fiber



B. A7 EGF



C. A7 scFv C242



D. A7 scFv G250



E. A7 Z_{IgG}



F. A7 Affi IgA



G. A7 Z_{IgG}/Z_{IgA}



H. A7 Z_{IgG}/Z_{IgG}



I. A7 Z_{IgG} Xa Knob



(57) Abstract: The present invention describes a modified virus comprising one or more non-native polypeptides, which polypeptide comprises one or more framework moieties each containing one or more binding moieties, which polypeptide is capable of being expressed in the cytoplasm and nucleus of a mammalian host cell in a conformation which is maintained in the absence of a ligand for said binding moieties, said conformation allowing said binding moieties subsequently to bind with said ligand, and which polypeptide is capable of transport through the nuclear membrane, wherein said modified virus has an altered tropism conferred by said binding moieties and the use of such viruses in the therapy, particularly in the treatment of tumours or other cancerous cells.



LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Modified Virus

5 The present invention relates to novel recombinant
viruses suitable for use in gene therapy. Such
recombinant viruses exhibit an altered tropism conferred
by incorporation of one or more non-native polypeptides
into one or more viral components, or the replacement of
such components by non-native polypeptides. These non-
10 native polypeptides can comprise elements which
mimic the structure of the original viral component so
as to permit inclusion of the polypeptide in functional
virion particles. These non-native polypeptides can
also comprise elements which confer a non-native ligand
15 binding function (i.e. altered tropism) to such virion
particles. Essentially, the non-native polypeptides
employed in the invention have primary and secondary
amino acid structures that enable their correct folding
in the nucleus or cytosol of mammalian host cells and
20 their subsequent transport through the nuclear membrane.
Such structures permit the assembly of functional
recombinant virion particles with an intact non-native
binding function and thus an altered tropism.

25 Clinical gene therapy was introduced in 1989. The aim
at that time was to correct gene defects in the immune
system via the *in vitro* introduction of a healthy gene
into the defective cells of the patient and transfusion
of the treated cells back to the patient. Since that
30 time, the indications and possible molecular uses of
gene therapy have increased dramatically. Today, ten
years after its introduction, one can envisage the use
of gene therapy to treat e.g. vascular diseases, cancer,
inflammatory diseases and infectious diseases such as
35 HIV.

At present, however, gene therapy is still not a useful

- 2 -

method in human medicine. One main reason is that gene therapy demands the packaging of the genes to be delivered into gene-carriers, or vectors, which can be administered to patients and which will target the genes only to the intended cells. Such vectors have so far not been available in reliable form.

Several classes of viruses have been considered as vectors for gene therapy applications, the most commonly used being adenoviruses, retroviruses, lentiviruses and adeno-associated viruses.

Ideal vectors for gene therapy would be those which can be administered systemically and yet deliver the desired genetic material specifically to desired cells or tissues. However, the viruses currently considered for human gene therapy applications have a broad tropism, being able to infect many different types of cells in the human body. This limits the potential safety of viral vectors and prohibits their use for systemic administration. For this reason, the development of targeted virus vectors, capable of infecting only selected cells, has been described as the holy grail of gene therapy.

One of the most widely investigated group of vectors for gene therapy applications is the adenoviruses. Human adenoviruses are divided into six hemagglutination groups (A-F) and each hemagglutination group is further subdivided into different serotypes. All in all there exist more than 40 different human adenovirus serotypes. The adenovirus which has been most frequently used for human gene therapy is adenovirus type 5 (Ad5) which belongs to hemagglutination group C.

Adenoviruses (Ad) are DNA viruses without an envelope, shaped as regular icosahedrons with a diameter of 60-85

- 3 -

nm.

The adenoviral capsid comprises 252 capsomeres, 240 hexons and 12 pentons (Ginsberg et al., Virology, 28, 782-83 (1966)). The hexons and pentons are derived from three viral proteins. The hexon comprises three identical proteins of 967 amino acids each, namely polypeptide II. The penton contains a base, which is bound to the capsid, and a fiber, which is non-covalently bound to, and projects from, the penton base. Proteins IX, VI and IIIa also are present in the adenoviral coat and are thought to stabilise the viral capsid.

Cell binding takes place through the fiber proteins, anchored to the virion at the vertices of the icosahedron. The fiber protein is not necessary for assembly and release of intact virions. Assembly of virions take place in the nucleus of infected cells.

The fiber protein, which is a homotrimer of a fiber polypeptide (namely adenoviral polypeptide IV), contains three functionally different parts: an N-terminal tail anchoring the fiber non-covalently to the penton base in the virion and which furthermore contains the nuclear-localization signal; a shaft domain comprising a variable number of repeats of a ~15 amino acid fiber shaft motif, (e.g. which is repeated six times in Ad3 and 22 times in Ad2 and Ad5); and a C-terminal globular domain, the knob, which contains the ligand which binds to the cellular Ad-receptor (Chrobozek et al., Microbiology and Immunology, (1995), p. 163-200). The knob is also functionally responsible for fiber trimerisation (i.e. it incorporates a trimerisation motif).

Each shaft repeat has two three-amino acid regions which

- 4 -

form beta-sheets and two amino acid regions which constitute the turns of the native extended fiber shaft. The crystal structure of the trimerised, cell binding domain has been determined and shows a unique topology different from other anti-parallel β -sandwiches (Xia et al., Structure 2: 1259-1270, (1993)). Binding of the fiber to the penton base of the virion can take place also in a cell-free system, i.e. the fiber can bind to fiberless virions (Boudin et al., Virology, 116: 589-604, (1982)).

Efforts to modify adenovirus fibers (e.g. produce recombinant fiber proteins) in order to modify the properties of adenovirus vectors, for example by altering tropism or cell binding, have been made and have been reported in the prior art.

The adenovirus fiber protein performs several biological functions which must be retained in order to produce active virus particles. The following fiber features are deemed to be of key importance in the construction of functional modified fiber proteins:

- i) the ability to form parallel homotrimers. This function is mediated by the N-terminal amino acid sequence of the wild type fiber knob.
- ii) the ability to bind to the penton base to form penton capsomeres. This function is mediated by the wild type fiber tail.
- iii) the ability to express a cell binding ligand allowing for attachment to target cells. In the native situation, this function is mediated by the wild type fiber knob (which binds to the cellular Ad-receptor).
- iv) the capability of transport into the nucleus of infected cells, which is vital to virus formation. This function is mainly, but perhaps not exclusively, mediated by the wild type fiber tail.

- 5 -

Previous attempts to change the tropism of adenovirus have involved genetic modification of fibers and knobs. However, this approach has not proved to be very successful. A major problem has been the incorporation of novel ligands in a functional context, which are capable of changing the tropism without interfering with the trimerisation of the fiber. For example, a short peptide ligand has been added C-terminally of the knob (Michael et al., Gene Therapy, 2: 660-8, (1995)) and a nonapeptide has been introduced into one of the knob "loops" (Dmitriev et al., J. Virol., 72(12):9706-9713 (1998)). However, the knob has a very complex structure due to interactions between the three fiber subunits which are necessary to conserve cell binding and trimerisation. Therefore, the knob only tolerates insertion of a few amino acids and no general method for construction of functional, genetically re-targeted adenovirus fibers exists to date.

Attempts have also been made to introduce new ligands into other parts of the adenovirion. By introducing the FLAG tetra-amino acid motif into the Ad penton, it has been shown possible to target Ad to cells normally not infected by Ad. The re-targeting was achieved by targeting with a bi-specific antibody where one specificity was directed against the FLAG peptide and the other against the new target cell (Wickham et al., J. Virol., 70: 6831-6838, (1996)).

A previously unaddressed problem encountered in the production of efficient recombinant viruses for gene therapy, is that of ensuring the functional folding of recombinant components upon expression in the nucleus and cytosol of host cells. This is particularly relevant where the wild-type virus to be engineered employs component expression in these intracellular locations during replication, for example, as in

- 6 -

adenovirus. Folded protein structures, particularly those which rely on disulphide bonds in cysteine bridges to maintain a functionally correct conformation, for example those derived from antibodies, can be rendered mis-folded and non-functional when expressed in the reducing environment of the nucleus and cytosol as part of a recombinant viral component. Thus, there exists in the art an unsatisfied and previously unappreciated need for protein structures which can retain a functional conformation when expressed as viral components in these cellular compartments, such that the binding functions which the structures comprise or support retain a binding function, particularly in the absence of the ligand for which binding is desired.

There are therefore major problems associated with the genetic engineering of Adenovirus fibers useful in the construction of recombinant re-targeted adenovirus (Ad-virus) for human gene therapy. These problems are deemed to be particularly important since more patients have been treated with adenovirus vectors than with any other type of vector (Trapnell et al., Biotechnology, 5: 617-625, (1994)). The present invention is directed towards circumventing such problems in the construction of genetically re-targeted viruses for gene therapy, where the new viral tropism has been accomplished by the introduction of a new cell binding ligand into a viral component protein.

The invention described herein relates in part to the production of functional recombinant adenoviral fiber proteins with a new tropism facilitated by removing or ablating (e.g. blocking or inactivating) the native cell binding domain and either replacing it with, or adding, a non-native polypeptide comprising an external cell binding ligand and an external trimerisation motif.

- 7 -

Curriel et al. in WO 99/41359, Wickham et al. in WO 98/54346 and Spooner and Epenetos in US 5,885,808 disclose various modifications to adenoviral components with the aim of producing recombinant viruses with an altered tropism. None of these disclosures provide a solution to, or indeed address or even recognise, the problems associated with the expression of functional non-native viral components in the nucleus and cytosol of host cells, a problem solved by the present invention.

Thus, in one aspect the invention provides a modified virus comprising a non-native polypeptide, which polypeptide comprises one or more framework moieties each containing one or more binding moieties, which polypeptide is capable of being expressed in the cytoplasm and nucleus of a mammalian host cell in a conformation which is maintained in the absence of a ligand for said binding moieties, said conformation allowing said binding moieties subsequently to bind with said ligand, and which polypeptide is capable of transport through the nuclear membrane, wherein said modified virus has an altered tropism conferred by said binding moieties.

A "modified" virus according to the invention is thus a virus which differs to a native (i.e. wild-type) virus. Generally speaking, the virus is modified such that a component of the virus is altered structurally over a native or wild-type (i.e. unmodified) component, or such that a structural component or feature is added to the virus, which is not present in the native or wild-type form. Advantageously, according to the present invention, a property or behaviour of the component is altered. In other words, the modified virus, differs functionally over an unmodified (native/wild-type) virus (e.g. by exhibiting an altered tropism). As discussed

- 8 -

above (and also further below), the virus is conveniently modified using genetic engineering techniques. Accordingly, a modified virus according to the present invention is advantageously a recombinant virus.

A modified virus according to the invention can be derived from any virus, and in particular any virus which may be used as the basis of a viral vector for gene therapy. Representative viral families include adenoviruses, retroviruses, lentiviruses and adeno-associated viruses and particularly include members of the family Adenoviridae or other virus families where viral structural components are synthesized and/or assembled in the nucleus or cytosol of the host cell, such as Reoviridae, Picornaviridae, Parvoviridae, Papovaviridae and Caliciviridae. In a preferred aspect, modified viruses of the invention are modified forms of adenoviruses, in particular Human adenoviruses and more particularly Human adenovirus type 5.

"Non-native polypeptide" as defined herein is a polypeptide sequence of two or more amino acids, the complete sequence of which is not found in a functionally equivalent position in the amino acid sequence of the wild-type virus, or more particularly in the wild-type viral component protein to be engineered.

In certain preferred embodiments of the invention, the "non-native polypeptide" may also be non-native in the sense of not occurring in nature, i.e. being a synthetically or artificially constructed or prepared polypeptide.

According to the present invention, the conformation of the non-native polypeptide after expression in the intracellular environment and, more importantly, also

- 9 -

its conformation in the extracellular environment, should be such that it is capable of binding to a desired ligand, e.g. a receptor or other molecule expressed on the outer surface of a target cell for the
5 modified virus.

As mentioned above, advantageously according to the present invention, the non-native polypeptide, when expressed in the cytoplasm of a mammalian cell has a
10 conformation which may be maintained in the absence of a ligand for the binding moiety(ies) of the polypeptide. Such a conformation is thus a stable conformation. In other words, upon expression in the cytoplasm, the polypeptide assumes a conformation which is maintained
15 in the cytoplasm, and when the polypeptide is transported into the nucleus and incorporated or assembled into a viral particle. Moreover, the conformation is such that the binding moiety or moieties of the polypeptide is or are able to bind to their
20 ligand, when exposed to it.

Thus, an important feature of the non-native polypeptide of the invention is that it is able to fold correctly in the cytoplasm or cytosol of a mammalian cell i.e. assume
25 a tertiary or three-dimensional structure which permits the functionality of the binding moiety(ies) to be retained (i.e. that the binding moieties retain their binding activity towards their ligand). In other words, the binding moiety remains capable of functional binding
30 to its ligand.

The solubility of recombinant proteins is considered a good indicator of their correct folding. Therefore non-native polypeptides of the invention may advantageously
35 be chosen not only on the functionality of binding moieties, stability of conformation and ability to form part of an assembled virion, but also on the basis of

- 10 -

solubility within cells in which a non-native polypeptide, or a viral component protein containing or comprising a non-native polypeptide, are expressed. For example, recombinant fiber proteins may be tested for their total expression in eukaryotic (e.g. insect) cells and the proportion recovered in the soluble fraction of the cell lysates determined.

The Inventors have found that phenotypic analysis of fiber-ligand fusion constructs expressed as recombinant proteins in baculovirus-infected insect cells showed that their degree of solubility correlated with the recovery of viable recombinant adenovirus. Therefore, it is preferred that any non-native polypeptide of the invention, or a viral component protein containing or comprising a non-native polypeptide, for example a fiber-ligand fusion construct, is characterised with regard to its solubility and target cell attachment as recombinant protein, prior to the re-insertion of the corresponding gene into the viral, for example adenovirus genome.

Thus the invention also provides a modified virus wherein said non-native polypeptide is selected using a solubility assay of the non-native polypeptide or the viral component protein comprising the non-native polypeptide. The non-native polypeptide is selected such that greater than 25%, preferably greater than 30%, more preferably greater than 50% or even 70%, of the non-native polypeptide or the viral component protein comprising the non-native polypeptide is present in the soluble fraction of cell lysates of cells expressing the non-native polypeptide or viral component protein comprising the non-native polypeptide. A suitable assay is described in the examples herein which may be applied *mutatis mutandis* to other cell expression systems. Reference herein to a non-native polypeptide or a viral

- 11 -

component comprising said non-native polypeptide being soluble in the cellular environment should be interpreted in line with these limits, i.e. when at least 25-30% of the polypeptide is present in the soluble fraction of a cell lysate, the polypeptide can be considered 'soluble'.

Thus, according to a further aspect, the present invention provides a method of determining the suitability of a non-native polypeptide (e.g. a recombinant viral fusion protein) for use in the preparation of a viral vector by determining its solubility in a cell system. More specifically is provided a method of assaying the solubility of a non-native polypeptide or a modified viral protein component of the invention, comprising the steps of i) expressing said non-native polypeptide or a viral component protein comprising said non-native polypeptide in permissive cells; ii) subjecting the cells to lysis to produce a cell lysate; iii) separating the soluble and insoluble fractions of the cell lysate; iv) analysing the soluble and insoluble fractions of the cell lysate for the content of said non-native polypeptide or viral component protein comprising said non-native polypeptide; and, v) comparing the relative content of said non-native polypeptide or viral component protein comprising said non-native polypeptide in the soluble and insoluble fractions to determine the solubility.

A ligand for a binding moiety of the invention may be regarded as the ligand which corresponds to the binding moiety, or in other words, the ligand to which the binding moiety was designed or selected to bind. The ligand is thus capable of binding to the binding moiety. The ligand and its binding moiety may thus be regarded as members of an affinity-binding pair, the ligand being

- 12 -

a binding partner for the binding moiety.

5 The binding moiety thus has a binding specificity for, or is capable of binding specifically to, a desired ligand. By "binding specificity" or "binding specifically" it is meant that the binding moiety is capable of binding to the desired (or "target") ligand in a manner which is distinguished from the binding to non-target molecules or ligands. Thus, the binding moiety either does not bind to non-target molecules or exhibits negligible or substantially-reduced (as compared to the target ligand) e.g. background, binding to non-target molecules. The binding moiety thus specifically recognises the target ligand.

15 This specificity of binding of the binding moiety thus permits the modified virus to be selectively targeted. In other words, the binding moiety may be designed or selected to enable the modified virus to bind to a desired or "target" cell. A binding moiety may be designed or selected which binds to a ligand expressed by or on a target cell, e.g. on the cell surface.

20 The ligand may thus be any desired ligand, and advantageously will be a molecule expressed on the surface of a target cell, in which it is desired to achieve expression of the modified virus. The ligand may thus be a cell surface receptor, or a cell-surface antigen. The ligand may conveniently be a protein or polypeptide molecule, but may be of any molecular nature, for example a lipid or carbohydrate.

25 In this manner, the modified virus may be modified to bind to a desired target cell to which the native (wild-type) virus from which it is derived does not bind, or it may be modified to have a more restricted binding specificity than the wild-type virus, in other words, to

- 13 -

bind to only a selected or particular sub-set or type of target cell from among a broader population of target cell types to which the wild-type virus binds. The tropism of the virus is thus altered. Hence, by
5 "altered tropism" it is meant that the modified virus exhibits a target cell binding specificity which is altered, or different, to that of the wild-type virus from which it is derived.

10 Generally, a non-native polypeptide according to the invention comprises at least one framework moiety and one or more binding moieties and is capable of interacting with other viral components to form a functional and infective virion particle. The ability
15 to form part of a functional virion particle can be facilitated by the presence of one or more sequences within the non-native polypeptide that effect binding to other viral components by the non-native polypeptide itself, or a viral component in which the non-native
20 polypeptide is comprised, in the virion assembly process.

A non-native polypeptide of the invention may replace or be incorporated into any viral component protein which
25 is capable of interacting with a target cell. The modified (e.g. recombinant) viral component possesses a cell binding function either by the nature of a retained native structure of the viral component itself or, by a new structure conferred upon it by the incorporation of
30 the non-native polypeptide and any structures comprised therein. In a preferred aspect of the invention, the non-native polypeptide is introduced or incorporated into or forms a fusion protein with, a viral protein component of the wild type virus, for example, an
35 adenoviral fiber protein, and especially preferably it is incorporated such that the wild-type fiber knob (or at least the cell-binding domain thereof) is removed

- 14 -

(i.e. replaced).

However, in an alternative embodiment, a wild-type fiber knob (or cell-binding domain thereof) may be retained, and a further, or additional, cell-binding domain may be added by virtue of the binding moiety(ies) of the non-native polypeptide. It is a feature of the invention that the cell binding functions are altered from those of the wild type virus to be engineered, such that an altered viral tropism is present in the modified virus. Where the wild-type fiber knob (or cell-binding domain thereof) is retained, this altered tropism may be achieved (or facilitated) by modifying the wild-type knob/cell-binding domain thereof to inactivate or block the native or wild-type cell binding function.

As will be discussed further below, it may in certain circumstances be desirable to construct or engineer a modified virus in which control (e.g. temporal control) may be exhibited on the expression of the altered tropism. Thus, in such a modified virus, a native or wild-type tropism may be retained, by retaining the wild-type fiber knob or cell-binding domain thereof, in addition to providing the non-native polypeptide which confers altered tropism. For example, this is desirable for the propagation of a modified virus in conventional cell lines known in the art via the additional presence in the modified virus of a wild type binding function, for example through the controlled expression of a wild type adenoviral fiber gene from an inducible control element (e.g. promoter).

The wild-type tropism may be ablated when desired by controlling (e.g. preventing) expression of the additional wild type viral component genes, or after expression, by removing or inactivating the wild type knob/domain, in such a manner that altered tropism conferred by the non-native polypeptide may be

- 15 -

expressed. An "altered tropism" according to the invention thus includes a "potential" altered tropism, i.e. the potential to express an altered tropism. It also includes an altered tropism which is additional to a wild-type tropism.

As mentioned above, the modified virus of the invention is preferably prepared using genetic engineering techniques and in preferred embodiments of the invention, the non-native polypeptide is provided as part of a fusion protein with a viral protein component, preferably as a fusion protein with an adenoviral fiber protein. Such a fusion protein, or more generally, such a modified viral component protein, represents a separate aspect of the present invention. In a preferred embodiment of this aspect of the invention, the modified viral component is a modified adenoviral fiber protein comprising a non-native polypeptide as defined above.

Techniques for preparing such non-native polypeptides and introducing them into viruses or viral components are well known in the art and widely described in the literature. Thus, for example, molecular biology or genetic engineering techniques are readily available, to prepare or construct genetic sequences capable of being expressed as a modified virus, or viral component, according to the present invention. As described further in the Examples below, a nucleic acid molecule or nucleotide sequence encoding a viral component protein, such as the adenoviral fiber protein, may be modified so as to introduce a nucleotide sequence encoding the non-native polypeptide, for example so as to encode a fusion protein comprising all or part of an adenoviral fiber protein and the non-native polypeptide.

Depending upon the viral component protein into which

- 16 -

the non-native polypeptide of the invention is incorporated, or which component(s) it replaces, a non-native polypeptide according to the invention can optionally comprise a further element which mimics the native structure or function of the viral component (i.e. is a functional equivalent of the viral component) so as to facilitate the assembly of functional virion particles. In the case of modified adenoviral fibers according to the invention, the functional integrity of the adenoviral fiber with regard to virion assembly, and particularly capsid assembly, is maintained by the presence of external amino acid trimerisation motifs such as the helical amino acid motif derived from the neck region of human lung surfactant protein D (Hoppe et al., FEBS Letters, 344: 191-195 (1994)). This and other trimerisation motifs known in the art can be functionally engineered into the fiber shaft and act as a fiber trimerisation signal to create knob-less fibers. For example, trimerisation motifs suitable for inclusion in modified viruses are described in WO 98/54346 and WO 99/41359. In a preferred embodiment of the invention, the non-native trimerisation motif present in the non-native polypeptide is the neck region peptide from human lung surfactant D.

As described in the Examples below, it may be convenient or necessary for any such additional or external motif or feature to be incorporated into the virus by means of a "linker" sequence. Such construction techniques for incorporation of DNA or amino acid sequences, via attachment to a linker sequence are known in the art and are within the routine skill of a protein/genetic engineer.

A "framework moiety" as defined herein, is a polypeptide (e.g. a protein) structure which retains a functional (e.g. folded) structure (or conformation) in the nuclear

- 17 -

and cytosolic cellular environments. Such functional structures are structures which allow a binding moiety attached to, or incorporated within, the framework moiety to retain ligand binding conformation in the absence of the ligand. This facilitates subsequent binding to that ligand, for instance, once the reassembled virion has left the cellular environment.

A framework moiety may thus be regarded as a type of "molecular scaffold" structure, which provides a framework to support or "hold" the binding moiety in an appropriate presentation or conformation to permit binding to its ligand. The framework moiety also provides the intramolecular interactions making a stable conformation in the cytosol possible. The framework moiety may thus be a protein or polypeptide molecule, which assumes a particular conformation or structure, and which tolerates modification in a particular region or regions, for example modification by amino acid sequence addition, insertion, deletion or substitution, or indeed insertion of a polypeptide sequence.

A "binding moiety" as defined herein, is thus a polypeptide structure attached to or comprising part of a framework moiety of the invention and which retains a binding function for a desired (target) ligand after expression of the non-native polypeptide of which it is part in the nucleus and cytosol of a host cell.

The binding moiety may be a contiguous or non-contiguous sequence of amino acids, and may be viewed as providing a binding site for the target ligand. A binding moiety may thus be provided by a region of a protein or polypeptide molecule, for example a surface region, e.g. a number of "surface" amino acid residues.

In certain preferred embodiments of the invention, the

- 18 -

framework and binding moieties may be of different origin, e.g. obtained from, or derived, from different sources, (e.g. from different proteins), for example by "grafting" or linking a desired binding moiety onto or to a desired framework. In other preferred embodiments, a binding moiety may be created within a protein "framework" structure, by modifying certain regions or residues of the framework protein, as described further below. In such an embodiment, the framework moiety may be represented by the "constant" or unmodified residues, and the binding moiety by the "variable" or modified residues.

A non-native polypeptide according to the present invention may thus be a combinatorial protein, that is a protein made by randomisation (random mutagenesis) of a particular protein structure, to generate a binding protein with novel, modified or enhanced binding characteristics. Such synthetically constructed "artificial" (in the sense of non-native) proteinaceous affinity binding molecules (i.e. proteins engineered to possess a particular or novel binding function) are known generally in the art.

Such combinatorial proteins can be prepared using various peptides and proteins as starting structures (Nygren and Uhlén, Current Opinion in Structural Biology, 7:463-469, 1997). Such proteins are known in the art, and may typically be prepared by random mutagenesis of a target protein, expression of the full library of these variants, e.g. on the surface of filamentous bacteriophage, followed by selection of a protein exhibiting the desired binding characteristics. This selection may typically involve a binding reaction between the variant protein and a target ligand (binding partner), which conveniently may be immobilised, and may be carried out *in vivo* or *in vitro*. The mutagenesis is

- 19 -

random in that the resulting amino acid encoded by any particular codon is not generally pre-determined but the positions where mutations are to be introduced are generally identified in advance. The mutagenesis may
5 involve amino acid substitution, deletion, or addition (e.g. insertion).

Preferred combinatorial proteins for use as non-native polypeptides of the invention are proteins known
10 generally in the art as affibodies. The term "affibody" as used herein defines an affinity binding molecule which is derived from a bacterial receptor protein, or binding domain thereof, wherein the binding domain is modified (e.g. by protein/genetic engineering) to modify
15 (e.g. alter or enhance) the binding properties thereof. Advantageously, the affibody is a non-native protein (in the sense of not occurring in nature) and is further preferred to have a novel binding site. Examples and further descriptions of such protein molecules are given
20 in WO 95/19374.

The use of an expression system such as surface display on phage provides a crucial link between genotype and phenotype; there is a self-contained unit which can be
25 selected on the basis of its specific binding interactions and which also carries the nucleic acid encoding for the protein responsible for the observed binding characteristics. This enables expression in useful amounts of the protein selected for its binding
30 characteristics, such expression typically taking place in a transformed bacterial host.

The protein, selected by its ability to bind to a target ligand (e.g. a desired cell surface molecule) may then
35 be used to prepare the modified virus or viral component of the present invention, or more particularly a nucleotide sequence encoding the desired combinatorial

- 20 -

protein may be so used.

5 Techniques for construction of a combinatorial library of protein molecules and subsequent selection to obtain proteinaceous binding molecules having desired binding characteristics are known in the art (Nygren, P. and Uhlén, M. Current Opinion in Structural Biology (1997) 7: 463-469). Generally, a protein molecule, perhaps having intrinsic beneficial properties such as
10 temperature or pH insensitivity, or conformational stability, is used as a "scaffold" or "framework" and a combinatorial library is then constructed via random but targeted amino acid substitutions (or other mutations) of that protein molecule, in order to produce a library
15 of molecules having different binding characteristics. Surface residues are generally targeted for random mutagenesis.

20 In addition to phage display technology (Smith et al., Meth. Enzym. 217, 228-57, (1993)), other methods for library construction and selection include, for example, ribosomal display (Hanes et al., Proc. Natl., Acad. Sci. USA 94: 4937-4942 (1997)), peptides-on-plasmids (Schatz et al., Methods Enzymol., (1996) 267: 83-109), RNA-
25 protein fusion (Roberts et al., Proc. Natl. Acad. Sci. USA 94: 12297-12302 (1997)) and DNA-protein linkage (STABLE) (Doi et al., FEBS Lett, 457(2): 227-30, (1999)).

30 Suitable protein frameworks may simply be linear peptides but preferably the framework will possess a folded three dimensional structure which has the potential for higher affinities and is less susceptible to proteolytic degradation. Although a framework may be designed *de novo*, naturally existing proteins or domains
35 are usually selected for further engineering. For the avoidance of doubt, it is to be noted that throughout this specification the word "protein" is used to refer

- 21 -

to whole protein molecules as well as domains or fragments thereof, polypeptides or peptides.

5 The choice of protein framework depends on several parameters including an ability to be effectively expressed in a desired host cell (e.g a mammalian cell). The protein should also comprise sufficiently large regions on its surface which are tolerant to substitution (or insertion or deletion etc.) without
10 losing the overall three dimensional structure. If the library is to be produced synthetically, a small overall size is a prerequisite. Where the selected framework protein has a binding function, amino acid residues involved in that interaction may be a target for
15 randomisation. Randomisation may be performed in order to enhance known binding properties or to develop binding molecules with new specificities.

20 Suitable framework molecules are discussed in Nygren et al. (supra) and include cyclic peptides in a constrained sequence (the number of amino acid residues in such a constrained sequence is not critical and can be 5 or more, e.g. 5 to 10 or more, e.g. 40 or more), immunoglobulin-like scaffolds including Fv or single-
25 chain (scFv) domains, bacterial receptors such as the 58-residue one-domain *Staphylococcal* protein A (SPA) analogue Z (the "Z Domain" being a derivative of the B domain of SPA), or other domains or analogues of SPA, DNA-binding proteins particularly zinc fingers and
30 protease inhibitors.

Of particular interest is the bacterial receptor domain Z. Nord et al., in Protein Engineering 8(6):601-608 (1995), describe a method of constructing a
35 combinatorial library of protein molecules based on the Z domain, which can be applied to a range of framework molecules. The method described is solid-phase-assisted

- 22 -

and based on the stepwise assembly of randomised single-stranded oligonucleotides.

5 As an alternative to modifying amino acid sequences or residues within a molecule to create a binding moiety, a binding moiety may be introduced to a framework moiety, for example by insertion or addition of a polypeptide constituting or comprising a binding moiety. The binding moiety may thus be attached to a framework
10 moiety. In such a case, a binding moiety may be provided by an affinity binding partner for the desired target ligand.

15 Preferred binding moieties of the invention may be derived from, but are not limited to, ligands (i.e. binding partners) for cell surface receptors, anti-receptor antibodies (or antibody fragments or derivatives), cell specific peptides, single chain antibodies (ScFv), single domain antibodies, and minimal
20 recognition units of antibodies such as a complementary-determining regions (CDRs) of Fv fragments. A binding moiety may thus be obtained or derived from the antigen-binding site or antigen binding or recognition/region(s) of an antibody, and such an antibody may be natural or
25 synthetic.

Also envisaged within the scope of the invention are binding moieties derived from peptides or polypeptides in any form, including hormones, antibodies, T cell
30 receptors, affibodies and ligands identified from various protein libraries.

As mentioned above, an important feature of the non-native polypeptide is that conformation is maintained in
35 the cytoplasm and nucleus of a mammalian cell, such that the binding function of the binding moiety is retained. As further mentioned above, this is achieved by

- 23 -

providing a non-native polypeptide which maintains correct folding in the cytoplasm and nucleus of a mammalian cell. It has been found that such correct folding may be achieved using a non-native polypeptide which does not rely on disulphide bonding for conformation (i.e. which does not contain di-sulphide bonds). As will be discussed in more detail below, a further preferred feature of a non-native polypeptide according to the present invention is the presence of a α -helical structure.

WO 95/19374 describes several framework proteins, which share the beneficial feature of not relying on S-S bonds for their conformation. Such framework proteins are useful as framework moieties according to the invention. These include domains of bacterial receptors such as staphylococcal protein A ($\alpha\alpha\alpha$ type), protein G (IgG binding parts, $\beta\beta\alpha\beta\beta$ type), protein L ($\beta\beta\alpha\beta\beta$ type), and protein G (HSA binding parts, $\alpha\alpha\alpha$ type). Framework moieties according to the invention can also be derived from various bacterial receptors, for example, but not limited to those listed in the table 1:

Table 1: Examples of G⁺ bacterial receptors

Receptor[ligand] ^a	Origin
Fc[IgG]receptor type I	<i>Staphylococcus aureus</i>
type II	<i>Staphylococcus pyogenes</i> [group A]
type III	<i>Streptococcus</i> group C, G, L
type IV	bovine group G streptococci
type V	<i>Streptococcus zooepidemicus</i> [group C]
type VI	<i>Streptococcus zooepidemicus</i> S212
Fibronectin receptor +	<i>S. aureus</i> , streptococci
M protein	<i>Streptococcus pyogenes</i> [group A]
Plasmin receptor	Streptococci group A
Collagen receptor	<i>S. aureus</i> , streptococci

- 24 -

	Fibrinogen receptor	streptococci groups A, C, G
	Protein L[6 light chains]	<i>Peptostreptococcus magnus</i>
	Protein H[human IgG]	<i>Streptococcus pyogenes</i> [group A]
	Protein B[human IgA, A1]	<i>Streptococcus agalactiae</i> [group B]
5	Protein Arp[human IgA]	streptococci group A
	Serum albumin receptor	streptococci groups A, C, G

^a Ligand is indicated when not obvious from receptor name

10 Framework moieties according to the invention are not restricted to structures from bacterial receptors. Also useful in the invention are other polypeptides comprising α -helical structures, often referred to as α -helical coiled coils, which are known from different

15 sources (Cohen et al., Science 263:488-489, (1994); Harbury et al., Science 262:1401-1407, (1993)). The coil making up the framework in these peptides consists of repeats of amino acid sequences containing characteristic positioned hydrophobic residues. The

20 structure of α -helical coiled coils is not dependent on intra- or intermolecular disulphide bridges for stability. Examples of α -helical coiled coils are the neck region peptide from human lung surfactant protein D, members of the spectrin superfamily, the leucine

25 zippers and parts of the hemagglutinin in influenza virus.

Particularly preferred as sources of non-native polypeptides according to the present invention are

30 members of the three-helix bundle family (e.g. as exemplified by the Z-domain of staphylococcal protein A).

Thus, in one embodiment, the non-native polypeptide

35 according to the invention comprises a framework moiety which is based on the structure of a domain derived from a bacterial receptor. Preferred structures of the

- 25 -

framework moieties of the invention are derived from or based upon the alpha-alpha-alpha ($\alpha\alpha\alpha$) -three-helix bundle or the beta-beta-alpha-beta-beta ($\beta\beta\alpha\beta\beta$) structure classes. Also preferred are structures based upon or derived from the α -helical coiled coil family, particularly a member of any of the 2-, 3- or 4-helical coiled coil families.

In an embodiment of the invention, it is envisaged that the binding moiety is present within one or more of the helical bundles and/or one or more of the loops connecting these bundles.

In a further preferred embodiment of the invention, the framework moiety is based on the structure of a domain derived from staphylococcal protein A, streptococcal protein G or *Peptostreptococcus magnus* protein L.

In a further particularly preferred embodiment of the invention, the framework moiety is a derivative of the immunoglobulin binding Z-domain from staphylococcal protein A (Nord et al., supra).

In such embodiments the binding moiety may be created by combinatorial protein engineering as discussed above. In other embodiments, the domain or protein selected (e.g. the Z-domain) may be used in wild-type form as the non-native polypeptide of the invention.

As mentioned above, through combinatorial protein engineering, e.g. targeted to surface-located residues of the Z-domain, libraries may be constructed from which novel variants (i.e. novel binding molecules) termed Z-domain affibodies, may be selected by binding the desired target ligand (Hansson et al., Immunotechnology 4: 237-52 (1999)).

- 26 -

Preferred Z-domain-based non-native polypeptides may comprise the following amino acid sequences:

5 VDNKFNKEXXXAXXEIXXLPNLNXXQXXAFIXSLXDDPSQSANLLAEAKKLNDAQAPK
[SEQ. ID. NO.: 45];

and

10 VDNKFNKEXXXAXXEIXXXXXXXXXXXQXXAFIXSLXDXXXXXSANLLAEAKKLNDAQAPK
[SEQ. ID. NO.: 46],

where X is any amino acid.

15 In the above polypeptides, the conserved (i.e. specified) amino acid residues may be regarded as constituting the framework moiety, and the variable residues X, as together providing the binding moiety.

20 Furthermore, framework moieties according to the invention need not be dependent on alpha-helices for their stability, provided that the required conformation for ligand binding by the binding moiety is retained after expression of the non-native polypeptide in the nucleus and cytosol of a host cell. Such frameworks for
25 example include frameworks derived from certain antibodies and their derivatives, known in the art, which do not require the presence of disulphide bridges to maintain structure and can therefore be expressed as part of a recombinant viral component in the nucleus or
30 cytosol of a host cell, whilst retaining a functional conformation and functioning as binding moieties for a desired ligand. Other antibody structures within the scope of the invention include antibody structures in which structurally relevant cysteine residues have been
35 replaced with alanine residues, and which retain the binding specificity of the antibody.

- 27 -

Additional framework moieties are also envisaged within the scope of the invention, for example, sequences derived from antibodies (e.g. monoclonal antibodies) and antibodies converted into a single-chain format allowing for the construction of genetic fusions to, or the modification of, viral components. In such a case where an antibody-derived framework moiety is used, then similarly as described above, a binding moiety may be created by protein/genetic engineering techniques (e.g. by modifying certain amino acid residues) or a binding moiety may be introduced (e.g. linked or grafted) to the antibody framework.

Antibody fragments conferring a desired binding activity can be used as binding moieties through the grafting of complementarity determining loops or regions (CDRs) of an antibody with a desired tropism into a framework moiety capable of productive folding, i.e. capable of retaining the binding specificity(ies) of attached or incorporated binding moieties, in the cytoplasm and subsequent transport into the cell nucleus where the virus assembly takes place. In other words, a binding moiety may be a CDR of an antibody.

Further, according to the invention, certain antibody structure frameworks can be used as framework moieties, for example in conjunction with antibody-based or derived binding moieties e.g. specificity determining loops (CDRs), resulting in the directed construction of antibodies suitable for construction of re-targeted viral components.

An antibody framework may be used as a framework moiety (for example to receive CDR loop(s) as binding moiety(ies)) according to the present invention, provided that it is capable of productive folding in the cytoplasm and subsequent transport into the cell

- 28 -

nucleus. Single chain Fv fragments from monoclonal antibodies without disulfide bonds have been produced (Proba et al., J. Mol. Biol., 275: 245-53, (1998)). Such ScFv antibodies provided they meet the functional criterion above, may be used on framework moieties according to the present invention. The binding moiety may be the binding site of the ScFv antibody itself, or it may be further engineered as discussed above.

Intracellular selection of functional antibodies from a polyclonal repertoire has also been achieved (Gargano et al., FEBS Letters 414: 537-40, (1997)). This selection technology may be useful in the identification of antibody fragments suitable for genetic retargeting of adenoviruses, namely suitable for use as framework, and/or binding moieties according to the invention.

Other antibody frameworks which may be used, include those based on camel antibodies which are naturally devoid of light chains, or certain VH regions derived from conventional antibodies.

An example of an antibody framework according to the invention that is capable of productive folding in the cytoplasm, is a particular anti β -galactosidase single chain Fv fragment. This single chain variable chain fragment (VK, linker, VH) reactive with β -galactosidase and capable of being expressed in the cytoplasm has been previously described (Martineau P and Betton J-M: J. Mol. Biol. 292, 921-929 (1999)). Thus, the invention also provides a modified virus comprising a framework moiety which comprises a sequence encoding this anti β -galactosidase single chain Fv fragment [SEQ ID. 47.] and functionally equivalent variants thereof.

The invention is not limited to framework moieties based upon antibody and affibody (e.g. receptor) structures,

- 29 -

but is extended to include other structures provided these are capable of retaining the binding specificity of attached or incorporated binding moieties when expressed in the cytosol/nucleus of infected cells.

5

Two or more non-native polypeptides, e.g. having different binding specificities, may be present in a modified virus of the invention, or a non-native polypeptide may incorporate two or more different binding moieties, or indeed two or more different framework/binding moiety constructs.

10

Accordingly, the invention provides a modified virus which comprises a first non-native polypeptide which binds a target cell and a second non-native polypeptide which binds a production cell or permissive cell.

15

Non-native polypeptides according to the invention may thus be present in recombinant viral components as comprising bi- or multi functional framework moieties (or framework/binding moiety constructs) constructed through genetic fusion between two or more different framework moieties (framework/binding moiety constructs). Use of such framework moieties (or framework/binding moiety constructs) can, for example confer infectivity of multiple cellular targets to the recombinant virus.

20

25

Also provided by the invention is a modified virus in which the non-native polypeptide according to the invention comprises a cleavage site positioned in a location that enables a binding moiety of the non-native polypeptide to be cleaved from the modified virus, for example preceding the binding moiety before the distal end of the fiber relative to the assembled virion. Examples of suitable cleavage sites are sites susceptible to a Factor Xa enzyme or a protease such as

30

35

- 30 -

the human rhinovirus 3C protease. If cleavage is carried out at the genetic level, for example, within a nucleic acid molecule comprising a part or all of a recombinant adenoviral genome, a cleavage site susceptible to an appropriate restriction enzyme or ribozyme may be used, for example in a particular cell population in which the virus is produced or targetted.

Non-native polypeptides according to the invention are capable of transport through the nuclear membrane. As is known from the art, this is an essential feature of viral components during replication in a host cell where expression of such components takes place in the cytosol, and assembly of such components takes place in the nucleus of the host cell e.g. in adenoviral replication.

The non-native polypeptide of the invention may be viewed as performing a dual role or function, firstly as providing a new binding domain (i.e. conferring altered tropism) and secondly as functioning as a viral component, namely a role as a functional part of a viral component (viral protein e.g. viral capsid protein). The non-native polypeptide may thus contribute to or function as a part of a viral protein. In other words, the non-native polypeptide may play a structural role or function, in the construction or assembly of a virus particle.

Cells comprising, or infected by, the modified virus of the invention also fall within the scope of the invention. Such cells may be of any origin provided that the cell type is capable of harbouring or propagating the virus. Generally, however, such cells will be mammalian cells. Such cells may be transfected with a membrane component (e.g. a membrane protein) that permits infection of the cell by the modified virus and

- 31 -

thus permits propagation of a virus which has a non-native tropism in a specific cell line, for example where propagation in conventional cell lines known in the art is prevented by way of the altered tropism of the virus.

Nucleic acid molecules encoding the non-native polypeptides, or modified viral component proteins or modified viruses of the invention are also envisaged. Vectors comprising such nucleic acid molecules, or the nucleotide sequence encoding the non-native polypeptides, or modified viral component proteins or modified virus of the invention, either for propagation of modified virus or further engineering of modified viruses are also included within the scope of the invention.

In a further aspect the invention provides a method for producing a modified virus according to the invention in cell culture. In one embodiment, the method comprises the steps of: genetically modifying a virus to produce a modified (e.g. recombinant) virus containing a non-native polypeptide, which polypeptide comprises one or more framework moieties each containing one or more binding moieties, which polypeptide is capable of being expressed in the cytoplasm and nucleus of a human host cell and there assuming and maintaining a conformation in the absence of a ligand for said binding moieties, which allow said binding moieties to subsequently bind with a said ligand and which polypeptide is capable of transport through the nuclear membrane, wherein said recombinant virus has an altered tropism conferred by said binding moieties; infecting permissive cells with such a virus; culturing said cells to produce the virus (e.g. at a sufficiently high titre), and, harvesting, and optionally, purifying the modified virus produced.

- 32 -

Ad vectors can be made replication competent or incompetent for permissive cells. For tumour therapy, replication competent Ad has the potential advantage that it can replicate and spread within the tumour (Miller et al., Gene Therapy, 3: 557-559). This may theoretically result in an increase of the chosen effector mechanism over that obtainable with replication incompetent vectors. Furthermore, infectious virus may contribute to an anti-tumour effect by cytopathogenic effects in infected cells as well as by evoking an anti-viral immune response which may harm infected cells.

It is desirable that a means of controlling the replication of competent modified virus is available in gene therapy applications.

Other means of controlling the replication of modified adenovirus are within the scope of the present invention. For example, the modified virus according to the invention may comprise a gene encoding a viral protein required for viral replication present under the control of an inducible promoter or genetic element. For example, adenovirus pre-terminal protein (pTP), may be present under the regulation of a tetracycline responsive transcription activator (trTA) such that the pTP is only expressed in the presence of doxycycline. Alternatively, the modified virus according to the invention may comprise modifications to the genome such that the virus only replicates in cells which have a defect in the DNA synthesis - apoptosis regulatory pathways.

Conveniently this may be achieved in a modified virus of the invention by introducing the further feature of a cleavage site upstream of the binding moiety, or positioned in any such location that enables the binding moiety to be cleaved from the modified virus, e.g.

- 33 -

between the fiber shaft and the binding moiety.

Thus in a further aspect, the invention also provides a method of regulating the replication of a modified virus comprising the steps of: constructing a modified virus such that a cleavage site (e.g. a site susceptible to enzymatic or chemical cleavage) is positioned between a binding moiety required for cell infection and the remainder of the recombinant viral component of which the binding moiety forms part, and, bringing said recombinant virus into contact with a cleavage agent or cleavage means (e.g. an enzyme, chemical or in the case of a photo-labile cleavage site, light) capable of cleaving said binding moiety from said viral component and thereby preventing the recombinant virus from undergoing further infection cycles.

Where the means of cleavage in the method is enzymatic, the cleaving enzyme can be encoded within the genome of the recombinant virus and can be inducible. In a preferred embodiment the cleavage site is cleaved by a Factor Xa enzyme. The use of other proteases known in the art is also envisaged, for example, the human rhinovirus 3C protease, which is available as a fusion protein with GST (Walker et al., Bio/Technology 12:601 (1994)). This protease is active at 4°C, and recognises and cleaves the sequence LEVLFQ // GP.

Non-native polypeptides of the invention can be selected from libraries after screening of such libraries for correct nuclear and cytosolic folding of the peptide and a desired binding function in a manner similar to phage display techniques as known in the art. Such libraries may consist of candidate peptides fused to wild type or modified (e.g. recombinant) viral components according to the invention, for example adenoviral fiber proteins. A fiber fusion library can be expressed on adenovirions

- 34 -

and used to select for correct cytosolic and nuclear conformation by observing those candidate fusions which facilitate replication.

5 Modified virus according to the invention may also be constructed with both wild type and modified viral components present, or encoded in the genome, and if desired, with each component gene under the control of different genetic control elements, for example
10 promoters. Thus, a modified virus according to the invention may comprise a modified viral component (i.e. modified to include or comprise a non-native polypeptide) and an equivalent or corresponding viral component which is unmodified, for example a wild-type
15 fiber, and a modified fiber.

For example, a recombinant adenovirus can be constructed with a wild type fiber and a modified or recombinant fiber (i.e. a modified fiber comprising or incorporating
20 a non-native polypeptide) e.g. a modified fiber derived from a 'fiber-candidate binding peptide' fusion library, which is expressed under control of a different promoter in an E1 deleted adenovirus. For example, the inventors have shown that a recombinant adenovirus fiber gene
25 under the CMV promoter can be cloned into the multiple cloning site of the shuttle vector pAdTrack and that viruses can be produced that express both the WT fiber and the recombinant fiber by homologous recombination using the Ad vector pAdEasy (He T-C, Zhou S, DaCosta LT,
30 Yu J, Kinzler KW and Vogelstein B: A simplified system for generating recombinant adenoviruses. Proc Natl Acad Sci, USA, 95:2509-2514, 1998).

It is also possible to have one or more fiber genes (for
35 example, a wild type gene and a modified gene) present in an adenoviral vector genome under the control of one or more inducible promoters, thus allowing each gene to

- 35 -

be switched on or off independently. For example, in a desired Ad construct, the wild type fiber gene, displaced from its normal position and its control by the major late promoter (MLP), can be cloned under control of the hormone inducible MMTV promoter, inducible by dexamethasone. This promoter allows for expression of the wild type fibre in the vector propagating cells. preferably, the modified fiber is cloned in the same Ad vector genome, for example, downstream from a TRE (tetracyclin-responsive element) sequence element that is switched on the in the desired cell line. In cells expressing the tTA protein (a transcriptional activator which binds to TRE), TRE will be activated in the absence of Tc (tetracyclin) or Dox (doxacyclin). Alternatively and more favourably, Ad vector constructs can be propagated in cells which express reverse tTA (rtTA) where TRE, and expression the modified fiber, is activated in the presence of Tc (or Dox).

The recombinant fiber proteins in the library can have the wild-type fiber knob replaced by an external trimerisation motif and one or more members of a peptide library (or other non-native polypeptide). Screening of the recombinant fiber proteins for a desired binding activity can be carried out in a manner analogous to phage display once the recombinant adenovirus has been propagated via expression of the wild type fiber in a culture of propagator cells.

Accordingly, modified virus with an altered tropism will require cells which it is capable of infecting in order to allows its propagation. Thus the invention also provides permissive cells for virus according to the invention which are capable of being cultured to propagate the virus. A replication incompetent virus (which generally comprises a deletion in its genome

- 36 -

rendering it incompetent) needs special producer or propagator cells which are able to supply the genetic information that is deleted or missing from the virus, in order to replicate. Such cells or cell lines are known in the art. A modified virus with an altered tropism may be unable to infect such cells as are known in the art. Accordingly the invention includes the modification of such a propagator cell to include a binding partner (ligand) for the binding moieties of the modified virus (e.g. such that the cells express such a ligand on their surface).

Accordingly the invention provides a cell, preferably an *in vitro* or *ex-vivo* eukaryotic cell, containing a modified virus or viral component of the invention.

It is also envisaged that modified virus with an altered tropism might be propagated in the same cells as the wild type virus by means of positioning the new tropism-altering binding moieties within a viral component which also comprises a wild type binding moiety (i.e. knob), with a cleavable site (e.g. for an enzyme such as Factor Xa or human rhinovirus 3C protease) between such moieties. After propagation in conventional cells via action of the wild type binding moiety, the viral component may be cleaved to remove the wild type binding moiety (e.g. the wild-type knob) and to reveal the new binding moiety and to confer the new tropism. It is also envisaged that the revealed binding moiety can be specific for a ligand which itself possesses a cell binding capability. For example, the binding moiety can be specific for the Fc region of an antibody, thereby permitting the use of an antibody which is cell specific for a desired cell type. Removal of the wild type binding moiety by cleavage and exposure of the virus to a cell specific antibody, allows the antibody to be adsorbed to the virus and thus targets the virus to the

- 37 -

cells specified by the antibody.

Non-native polypeptides according to the invention preferably confer an altered tropism to the recombinant virus in comparison with the tropism of the virus to which the non-native polypeptide has been introduced. Such altered tropisms allow recombinant viruses of the invention to be used in treatment of disease in human or animal subjects, either by *in vivo* treatment of, or *ex vivo* treatment of cells of, the subject requiring treatment.

The tropism of the modified virus of the invention may be altered such that the virus, via one or more binding moieties targets particular cells. Thus the invention also provides a modified virus wherein a non-native polypeptide comprises a binding moiety capable of binding to a cell specific ligand which may optionally be Prostate Specific Membrane Antigen, EGF receptor, Her-2/Neu, VEGF receptor, CD22, gp120, MHC/peptide complexes or membrane structures or surface molecules expressed or present on proliferating cells, tumor cells or virus infected cells.

There are many ways in which modified virus exhibiting an altered tropism according to the invention can be used in gene therapy applications. In the case of tumour diseases, the following options exist:

I. Use of vectors to introduce transgenes into tumours, such as:

- antisense oncogenes
- suicide genes
- genes for immune modulatory substances or tumour antigens
- genes for anti-angiogenic factors

- 38 -

5 Examples of transgenes suitable for inclusion in the modified virus of the invention are antisense oncogenes, suicide genes, genes for immune modulatory substances, genes for tumour antigens, genes for anti-angiogenic factors, cytokines, genes for vascular endothelial growth inhibitors, genes for fusogenic membrane glycoproteins, cytotoxic genes, a gene encoding an enzyme which converts a pro-drug to cytotoxic substance, a gene for cytosine deaminase, a gene for uracil
10 phosphoribosyl transferase. In a preferred embodiment of the invention the modified virus comprises transgenes encoding cytosine deaminase and uracil phosphoribosyl transferase either as separate genes or, more preferably, together as a bifunctional fusion gene
15

In methods of treatment employing viruses of the invention with the transgenes cytosine deaminase and uracil phosphoribosyl transferase, preferably the virus is present or is co-administrated with one or more
20 inhibitors of dihydropyrimidine dehydrogenase. This has the effect of increasing the toxic effect of the products formed by these transgenes from the pro-drug, 5-Fluorocytosine.

25 Also provided by the invention is a modified virus of comprising one or more viral components or transgenes present under the control of one or more inducible promoters or genetic elements, which are for example tissue specific or which respond to the presence of an
30 exogenous expression modulating substance.

35 II. Use of infectious virus. This has the added value over the use of non replicating vectors in that virus can spread from cell to cell within a tumour, thereby multiplying the initial hit, or effect, on the tumour. Tumour cell destruction can occur not only by the cell-destroying mechanism engineered into the vector but

- 39 -

also by the cell destruction which is associated with the virus infection per se and by the attack of the body's immune response on the virus infected cells. This principle has already been tested in man through the direct intra-tumoural injection of an adenovirus which has been made gene manipulated to replicate only in p53 mutant tumour cells. The experience from these limited trials on large "head-and-neck" tumours are partially encouraging with a total regress of 2/11 treated tumours which are otherwise resistant to any form of known treatment (Shen Y: Personal communication).

Also provided by the invention is a modified virus wherein the gene encoding adenovirus death protein (ADP) is placed under the control of a promoter permitting over-expression of the protein, such that the lytic capacity of said modified virus is increased.

Thus, for gene therapy the modified virus may further be modified, according to standard techniques, principles and proposals widely described in the literature, to incorporate or comprise a desired therapeutic gene or therapeutic nucleic acid molecule. "Therapeutic" is used broadly herein to include both therapy (in the sense of curative or palliative therapy of a pre-existing or diagnosed condition) and prophylaxis. The gene may encode a desired therapeutic product (e.g. a therapeutic polypeptide or antisense molecule).

In a further aspect of the invention, the modified virus may include a site (e.g. a restriction site) for insertion of a desired therapeutic gene/nucleic acid molecule.

One disadvantage of using a wild type virus in gene therapy, even after modification according to the

- 40 -

invention, is that the human or animal subjects to be treated with the virus may possess pre-formed antibodies to components of the wild type virus still present or only partially modified in the modified virus. For example human adenovirus serotype 5 hexon protein is often targetted by pre-formed antibodies. Whilst modification of a single viral component, for example the component containing the non-native polypeptide according to the invention, may be sufficient to reduce the initial immune response to the modified virus via pre-formed antibodies, it may be preferable to modify a further component of the virus to further reduce the immune response of the host.

Thus the invention further provides a modified virus according to the invention which further comprises a viral component which is replaced with an equivalent component or is modified such that binding of said virus by antibodies pre-formed to the wild type virus is reduced. For example, an adenovirus of serotype Ad5 may comprise a hexon protein which is swapped for a hexon protein of a different serotype against which pre-formed antibodies are present at a reduced level compared to serotype Ad5, for example the different serotype may be Ad37. Alternatively, a modified adenovirus may comprise epitope sequences of the hexon protein to which pre-formed antibodies bind which are modified to produce a recombinant hexon lacking immunogenic epitopes.

Furthermore, the modified adenovirus may comprise a hexon protein which further comprises a peptide capable of binding a protein, for example, a non-immune system protein such as human serum albumen, sufficiently to cover the immunogenic epitopes of the hexon which are bound by pre-formed antibodies.

Also provided by the invention is a modified virus

- 41 -

according to the invention for use in therapy or in the preparation of a medicament for the treatment of tumour cells or proliferating cells.

5 Additionally provided by the invention is a pharmaceutical composition comprising a modified virus of the invention and a pharmaceutically acceptable carrier or excipient.

10 The invention will now be described in more detail with reference to the following non-limiting Examples, in which:

15 Figure 1 shows a schematic description of the sequence of construction of different recombinant fibers.

A. The NRP sequence is supplied with flanking SphI and XhoI sites using PCR and ligated into WT fiber which has been supplied with flanking EcoRI and XhoI sites. The resulting fiber is called A1 and contains the fiber
20 tail, first shaft repeat and the NRP motif.

B. EGF is joined to the fiber A1 by SOE. In the process an amino acid linker and a ClaI restriction site is added between the NRP and EGF sequences. The resulting fiber is named A1 EGF. In this fiber, EGF can
25 be substituted for new ligands by ligation using the ClaI and XhoI sites.

C. The NRP-Linker-EGF part of fiber A1 EGF is subjected to PCR. In the process an upstream NheI site is introduced into the sequence in frame with the WT
30 fiber sequence. After AT cloning, the sequence is ligated into WT fiber using NheI and XhoI to create fiber A7 EGF. A7 differs from A1 in that A7 contains the first seven shaft repeats of the Ad5 fiber. In the A7 EGF construct, EGF can be substituted for new ligands
35 by ligation using the ClaI and XhoI sites.

Figure 2 shows a schematic representation of the

- 42 -

different recombinant fibers used in the present application.

- A. The Ad5 wild type fiber.
- B. The fiber A7 EGF described above.
- 5 C. The fiber A7 scFv C242 obtained by substituting EGF in fiber A7 EGF for the scFv fragment by ligation as mentioned above.
- D. The fiber A7 scFv G250 obtained by substituting EGF in fiber A7 EGF for the scFv fragment by ligation as
10 mentioned above.
- E. The fiber A7 Affi IgG where EGF in fiber A7 EGF has been substituted for an IgG binding affibody as mentioned above.
- F. The fiber A7 Affi IgA where EGF in fiber A7 EGF has
15 been substituted for an IgA binding affibody as mentioned above.
- G. G. The fiber A7 Affi IgG/Affi IgA where EGF in fiber A7 EGF has been substituted for an IgG binding affibody linked to an IgA binding affibody as mentioned
20 above.
- H. The fibre A7 ZIgG/ZIgG where EGF in fibre A7 EGF has been substituted for an IgG binding affibody linked to another IgG binding affibody as mentioned above.
- I. The fibre A7 ZIgG Xa Knob where EGF in fibre A7 EGF
25 has been substituted for an IgG binding affibody linked to a cleavable wild type fibre knob.

Figure 3 shows the binding of ZIgG affibody when incorporated into fibers expressed on virions.

30 Panel A. After binding of virus or protein to the membrane, it was incubated with Fc3(1) followed by HRP conjugated anti human IgG and developed. 1 = the virus A7 ZIgG Xa Knob; 2= the same virus after cleavage with Xa; 3 = WT virus; 4 = virus with two fibers e.g. WT and
35 A7 ZIgG; 5 = Fc3(1); 6 = Protein A; 7 = Protein AG.

Panel B. After binding of virus or protein to the membrane, it was incubated with Fc3 followed by HRP

- 43 -

conjugated anti human IgG and developed. 1 = the virus A7 ZIgG Xa Knob; 2= the same virus after cleavage with Xa; 3 = WT virus; 4 = virus with two fibers e.g. WT and A7 ZIgG; 5 = Fc3; 6 = Protein A; 7 = Protein AG.

5 Panel C. Control to show presence of virus on membrane. 1 = the virus A7 ZIgG Xa Knob; 2= the same virus after cleavage with Xa; 3 = WT virus. After binding of virus the membrane was incubated with HRP conjugated anti Ad5 hexon and developed.

10 Panel D. Treated as membranes in panel A and B but the incubation with Fc was omitted. 1 = the virus A7 ZIgG Xa Knob; 2 = WT virus; 3 = protein A; 4 = protein AG.

15 Explanatory note: Fc3(1) is known to bind ZIgG and protein A. Fc3 is known to bind protein G (AG) only.

Figure 4 shows a schematic representation of the fibers described in Example 11. The NRP-linkers were inserted in the *NheI* site upstream of NRP.

20 Figure 5 shows gel photos demonstrating expression and solubility of recombinant fibers in Sf9 cells. (A), whole cell lysate; (B), soluble fraction. Baculovirus-infected cells were harvested at 48 h after infection, lysed in isotonic buffer, and cell lysates divided in two aliquots. One aliquot was centrifuged at 10,000 x g for 10 min and supernatant was kept (panel B), whereas the other was analysed as whole cell lysate (panel A). Both aliquots were heat-denatured in 30 SDS-sample buffer, analysed by conventional SDS-PAGE, and blotted. Blots were reacted with 4D2.5 mAb (specific for the fiber tail) and radiolabeled secondary antibody. Immunoblots were quantitated by autoradiogram scanning. Quantitative data, expressed as the percentage of 35 soluble versus total fiber content, are shown in Table 7.

EXAMPLESGeneral procedures and starting materials.

Recombinant adenovirus fibers were constructed using methodologies based on ligation and PCR (Clackson et al., General application of PCR to gene cloning and manipulation, in PCR, A Practical Approach, Eds McPherson MJ, Quirke P and Taylor GR, IRL Press, Oxford, page 187, (1992)), i.e. PCR-ligation-PCR (Alvaro et al., BioTechniques 18: 746-750 (1995)) and splicing by overlap extension (SOE) (Horton et al., Recombination and mutagenesis of DNA sequences using PCR, in McPherson MJ (ed), Directed Mutagenesis, IRL Press 1991, p 217.). Gene products generated by PCR were generally cloned into the vector pCRII (Invitrogen Corp.) using so called TA cloning (Clark J.M., Nuc. Acids Res. 16: 9677-86, (1988)). Subclonings were performed according to standard methods (Sambrook et al., Molecular cloning. A laboratory manual. Second Edition. Cold Spring Harbor Laboratory Press, (1989)) in the vector pGEX-4T-3 (Amersham Pharmacia Biotech). Genes encoding recombinant fibers were sequenced using the Perkin Elmer ABI Prism sequencing equipment and were expressed in mammalian cells (SV40 transformed African Green monkey kidney cells, COS7, obtained from American Type Culture Collection, VA, USA) using vectors described below, and in insect cells (Sf9 cells from *S. frugiperda* obtained from American Type Culture Collection, VA, USA) using Baculovirus expression (Kitts et al., Biotechniques 14(5): 810-7, (1993)) (virus and vector from Clontech, Palo Alto, CA, USA) and stained with monoclonal antibodies specific for fiber tail, trimeric fiber and the new cell binding ligand. The following parameters were evaluated by immunostaining:

- i) nuclear transportation
- ii) functional expression of the new cell binding ligand

- 45 -

iii) ability to form trimers

Recombinant fibers were rescued into the Ad genome by a recently developed procedure (as described in Example 7 herein). The plasmid pTG3602 (Chartier et al., J. Virol., 70: 4805-4810, (1996)) containing the entire Ad5 genome as a PacI-PacI fragment was used as starting material. The approximately 9kb fragment of the genome between SpeI and PacI and containing the wild type fiber gene was cloned separately in pBluescript. From this fragment an approximately 3kb fragment between SacI and KpnI was further subcloned. A deletion of the native fiber gene with the exception of the N-terminal nucleotides upstream of the NdeI site of the fiber was created in the 3kb fragment and an XhoI site introduced in its stead allowing for ligation of recombinant fibers into the fiber-deleted 3kb fragment (the 3 kb fiber shuttle) between NdeI and XhoI.

The 3 kb fiber shuttle with recombinant fiber was re-introduced into the 9 kb fragment cut with NheI using homologous recombination in E.coli (Chartier et al., Supra). The resulting recombinant 9 kb fragment was finally excised from the vector with SpeI and PacI and joined to the isolated 27 kb fragment by Cosmid cloning.

The presence of an insert of the expected properties was verified in all cosmid clones by PCR. Cosmid clones were also restricted with Hind III and the presence of restriction fragments of the expected size verified on gels.

Recombinant Ad genomes were isolated after restriction with Pac I and used to transfect suitable cells. The occurrence of plaques was determined by microscopic inspection of the transfected cell cultures.

- 46 -

List of oligonucleotide primer names for the primers used (all given as 5'-3' sequence):

- [SEQ. ID. NO.: 20] Name 149: TTCCTCGAGTTATTCTTGGGCAATGTATGA
- [SEQ. ID. NO.: 21] Name 175: GGGGAATTCGATGAAGCGCGCAAGACCGTCTGAA
- 5 [SEQ. ID. NO.: 22] Name: 196: GCTCGAGTTATCCGTTTGGAAACAACCTCTAC
- [SEQ. ID. NO.: 23] Name: 228: CTCGAGTCATCTCAATTCCCACCACTT
- [SEQ. ID. NO.: 24] Name: 238: TGGCATGCCTGACGTAGCAAGCTTACGA
- [SEQ. ID. NO.: 25] Name: 253:
- GGGGAATTCATCGATGCAGGTCCAGTTGGTGCAGTCT
- 10 [SEQ. ID. NO.: 26] Name: 265: CAGGTCCAGTTGGTGCAGTCT
- [SEQ. ID. NO.: 27] Name: 269:
- GGGGGCCTGGGCGTCGTTTCAGCTTCTTGGCTCCGTTTGGAAACAACCTCTAC
- [SEQ. ID. NO.: 28] Name: 270:
- CTGAACGACGCCCAGGCCCCCAAGAGCGACCCATCGATCATGAACCTCCGACTCCGAATGT
- 15 [SEQ. ID. NO.: 29] Name: 273:
- CCCCTGGAGTTAAATTTTCTTGTCCACCTTGGTGCT
- [SEQ. ID. NO.: 30] Name: 274:
- GGGGAATTCATCGATGGACTACAAAGATATTGTGATGACGCAGGCT
- [SEQ. ID. NO.: 31] Name: 275: CTACCTCGAGTTAACACTCATTCCTGTTGAAGC
- 20 [SEQ. ID. NO.: 32] Name: 326: GGGGCTAGCCCCTGACGTAGCAAGCTTACGA
- [SEQ. ID. NO.: 33] Name: 403: GGG CTC GAG TTA CTC GAT GGG GGC TGG
- GAG GGC
- [SEQ. ID. NO.: 34] Name: 414:
- GGCCCCCGAGGCCTCGAGTGAGGAGACGGTGACCGTGGT
- 25 [SEQ. ID. NO.: 35] Name: 416:
- GGCCCAGCCCACGAATTCATCGATGGATATTGTGATGACGCAGGCT
- [SEQ. ID. NO.: 36] Name: 418: AGA CTG CAC CAA CTG GAC CTG
- (SNN)₁₈CCGTTTCAGCTCCAGCTTGGT (S is dA, dG or dC and N is dA, dG, dC or dT)
- 30 [SEQ. ID. NO.: 37] Name: 473: GGC AAT TCC ATC GAT CGC CAC CAT GGA
- CAT TGT GAT GAC CCA GTC T
- [SEQ. ID. NO.: 38] Name: 474: CCC CTC GAG TTA ACA CTC ATT CCT GTT
- GAA GCT
- [SEQ. ID. NO.: 39] Name: 476: ACC ACG GTC ACC GTC TCC TCA GCT GAT
- 35 GCT GCA CCA ACT GTA
- [SEQ. ID. NO.: 40] Name: 478: TGA GGA GAC GGT GAC CGT GGT
- [SEQ. ID. NO.: 41] Name: 503: GGGCCATCGATCGTAGACAACAAATTCACAAA

- 47 -

[SEQ. ID. NO.: 42] Name: 504: GGGCTCGAGTTATTTTCGGCGCCTGAGCATCATT

[SEQ. ID. NO.: 43] Name: 550:

TCGGTTTGGAAACAACCTCTACCTTTTTTTTCGGCGCCTGAGCATCATT

[SEQ. ID. NO.: 44] Name: 551:

5 AAAAAGGTAGAGTTGTTTCCAAACGGAGTAGACAACAAATTCAACAAA

EXAMPLE 1:

10 Genetic insertion of a trimerisation motif (the neck region peptide from human lung surfactant D) into adenovirus fibers.

The gene encoding Ad 5 WT fiber was obtained from a preparation of Ad5 virus by PCR using an upstream primer
15 (Primer 175) identical to the first six coding triplets of the fiber plus an EcoR1 site and a downstream primer (Primer 149) annealing to the six terminal coding triplets of the fiber plus an Xho1 site. The fiber thus obtained [SEQ. ID. NO.: 1] was cloned into the vector
20 pBluescript using these restriction sites and can be further sub-cloned into other vectors using the same restriction enzymes.

Fiber peptides were made where the knob was replaced
25 with an external trimerisation motif (see below). The purpose behind the introduction of an external trimerisation motif is two-fold: a) to remove the knob containing the native trimerisation signal but also the cell binding part of the fiber, and b) simultaneously to
30 supply the necessary trimerisation signal. In this case one particular amino acid motif have been used, i.e. the 36 aa "Neck Region Peptide" = NRP [SEQ. ID. NO.: 2] from human "Lung Surfactant Protein D" (Hoppe et al., Supra). It should be noted that the sequence used is slightly
35 longer than the actual trimerisation part of NRP in that the eight amino acids (KKVELFPN) following the trimerisation signal in human lung surfactant protein D

- 48 -

has been retained in all constructs containing the NRP sequence. The sequence KKVELFPN functions as an efficient linker between the trimerisation signal and the C-terminal carbohydrate domain of the human lung surfactant D and is considered to have the same important function in the recombinant fibers described herein. The DNA sequence coding for the trimerisation motif was synthesized, cloned and verified by sequencing.

10

In order to introduce the NRP motif into the adenovirus fiber, the NRP sequence was subjected to PCR (Clackson et al., Supra) with the upstream primer 238 containing SphI N-terminally of the NRP coding sequence and the downstream reverse primer 196 containing XhoI C-terminally of the coding sequence. After cutting with SphI and XhoI the NRP sequence was ligated into the WT fiber gene cut with the same enzyme. The resulting recombinant fiber A1 [SEQ. ID. NO.: 3] contains the fiber tail and the first shaft repeat followed by the NRP trimerisation motif. For a schematic representation of the constructions and construction pathways see Figs 1 and 2.

25

To replace the cell binding function of the knob a new cell binding ligand was subsequently introduced into the fiber in addition to the external trimerisation amino acid motif (see below).

30

EXAMPLE 2:

Assembly of gene construct encoding recombinant adenovirus fibers with epidermal growth factor (EGF) and the external trimerisation motif from human lung surfactant D.

35

For a schematic representation of the constructions and construction pathways see Figs 1 and 2.

- 49 -

The DNA sequence for human EGF [SEQ. ID. NO.: 4] was synthesized, cloned and sequenced in the project. This sequence was then joined to the A1 fiber mentioned above by splicing by overlap extension. In this case the EGF gene was subjected to PCR with an upstream primer (270) (identical to the first seven coding triplets) and containing an overhang with the sequence for an amino acid linker [SEQ. ID. NO.: 5] derived from Staphylococcal protein A and a ClaI restriction site, and a downstream primer (228) (complementary to the seven terminal triplets of the + strand and containing an XhoI site). The A1 fiber gene was subjected to PCR with an upstream primer identical to the first six coding triplets of the gene and an EcoRI site (175) and a downstream primer (269) complementary to the seven terminal coding triplets of the + strand and an overhang complementary to the overhang in Back primer for the EGF sequence. The two PCR products were then joined by PCR under standard SOE conditions (Horton et al., Supra) to produce fiber A1 EGF [SEQ. ID. NO.: 6].

In order to construct a fiber with the first seven shaft repeats, the NRP trimerisation signal, the Staphylococcal linker and EGF, the fiber A1 EGF was subjected to PCR with an upstream primer (326) identical to the first seven 5' triplets of the NRP sequence plus an upstream NheI site and a downstream primer (228) complementary to the seven terminal triplets of the A1 plus strand. After cloning the PCR product was restricted with NheI and XhoI and ligated into WT Fiber restricted with the same enzymes to obtain Fiber A7 EGF [SEQ. ID. NO.: 7] which is similar to A1 EGF but differs in that it contains the first seven shaft repeats of the Ad5 Fiber.

35

Example 3:Assembly of gene construct encoding recombinant

- 50 -

adenovirus fibers with single-chain antibodies and the external trimerisation motif from human lung surfactant D.

5 Two monoclonal antibody single chain fragments were used to construct recombinant adenovirus fibers. The first is a single chain fragment (scFv) of the monoclonal antibody G250, which with high selectivity has been shown to react with a protein antigen on human renal carcinoma cells (Oosterwijk et al., Int. J. Cancer 38: 10 489-94, (1986)). The second is a single chain fragment of the monoclonal antibody C242 which reacts with i.a. colorectal and pancreatic carcinomas (Johansson C., Thesis, University of G'teburg, (1991)).

15

G250 constructs

The single chain fragment (Variable kappa chain or VK, linker, variable heavy chain or VH, joining sequence and constant heavy domain 2 or CH2) of the antibody G250 was 20 constructed as previously described (Weijtens et al., J.Immunol., 152(2): 836-43, (1996)). This G250 construct is [SEQ. ID. NO.: 8]. To permit cloning into the aforementioned A1 and A7 fiber constructs, the single chain fragment was supplied with an upstream ClaI 25 site and a downstream XhoI site by PCR using primers 416 and 403.

C242 constructs

The single chain fragment of the antibody C242 (Variable 30 kappa, Linker, variable heavy and constant kappa or CK) [SEQ. ID. NO.: 9] was constructed as follows by SOE using cDNA from the antibody producing hybridoma as original templates. VKCK was amplified using primers 274 and 275, VHCH1 was amplified using primers 253 and 35 273. An scFv (single chain variable fragment) (VK LinkLib VH) was constructed by SOE as follows. VK and VH were amplified separately using primers 416/418 and

- 51 -

265/414 respectively with the above mentioned VKCK and VHCH1 as templates and joined together by SOE using primers 414 and 416. In the construct VK LinkLib VH, the linker between VK and VH is a randomized 18 amino acid sequence as described previously (Tang et al., J. Biol. Chem., 271: 15682-86, (1996)). The nucleotide sequence for this linker is present in primer 418. The construct VK LinkLib VH was cloned into the vector pAK100 (Krebber et al., J. Immunological Methods 201: 35-55, (1997)). Phage display and selection of antigen binders by panning was performed using methods described earlier (Krebber et al., Supra). In the present experiments, the CanAg antigen, reacting with antibody C242, was adsorbed onto biotinylated antibody C241 (which binds another epitope on the antigen than C242) bound to streptavidin coated tubes (CanAg Diagnostics Ltd, G'teburg, Sweden). Several binders were isolated and shown to contain different linkers. A particular VK Linker VH construct shown by sequencing to contain the linker PPDFVPPAASFDPHSPRG (one letter amino acid code) was selected for further work based on antigen binding ability. CK was linked to this construct by SOE to obtain the format C242 VK Link VHCK. In this SOE the VK LinkLib VH was amplified using primers 416 and 478 and the CK amplified with primers 476 and 474. The amplified products were then joined by SOE using primers 416 and 474.

In the PCR reactions mentioned above the gene sequences for the single chain fragments were supplied with an upstream Cla1 site (present in primers 416 and 473) and a downstream Xho1 (present in primers 403 and 474) to allow for ligation into the A1 and A7 fiber constructs mentioned earlier (for a schematic representation of the constructions and construction pathways see Figs 1 and 2) to construct the fibers A1 G250, A7 G250, A1 C242 and A7 C242 (Fig 1 and [SEQ I.D. NOS. 10-13]).

- 52 -

Example 4:Assembly of gene constructs encoding recombinant adenovirus fibers with affibodies and the external trimerisation motif from human lung surfactant D.

5

To investigate if mono and divalent antigen binding structures based on the structure of single staphylococcal protein A domains could be functionally expressed when incorporated into recombinant Adenovirus fiber, gene constructs for mammalian and insect cell expression were made. Assembly of gene constructs encoding recombinant adenovirus fibers containing the IgG binding Z domain (ZIgG) derived from staphylococcal protein A (Nilsson B., Prot. Eng. 1:, 107-13, (1987)) [SEQ. ID. NO.: 14] or a Z domain-derived IgA-specific affibody (ZIgA) selected using phage display (Gunneriusson E. et al., App. Env. Micro., 65: 4134-40, (1999)) [SEQ. ID. NO.: 15] was accomplished as follows.

20 The genes encoding the respective affinity moieties were amplified by PCR using primers 503 and 504 on the following plasmid templates; pEZZmp18 (Tang et al., Supra) for the Z domain construct and pKN1-dZiGA (Clackson et al., Supra) for the ZiGA construct.

25

In the PCR amplification, the genes for the two different affinity ligands were supplied with an upstream Cla I site and a downstream Xho I site in the appropriate reading frame for subsequent ligation into the above described fiber gene A7 EGF resulting in constructs encoding the recombinant fiber A7 ZiG [SEQ. ID. NO.: 16] and A7 ZiGA [SEQ. ID. NO.: 17], respectively, which were adapted for later being rescued into the Ad genome (see below) for the production of recombinant viruses carrying the new binding specificities. Furthermore, the fiber A1 ZiG was constructed by ligation of the modified ZiG into the

35

- 53 -

aforementioned fiber A1 EGF (see also Fig 1).

In addition, a third gene construct was assembled encoding a fiber containing both of the above mentioned affinity domains. This construct encodes a fiber containing the fiber tail, the first seven shaft repeats, the NRP sequence, the staphylococcal protein A linker, the IgG binding Z domain, the eight amino acid linker from NRP (KKVELFPN) followed by the IgA binding affibody. To construct this fiber the two different affinity domains were first genetically joined together by SOE using primers 550 and 551 with overhangs complementary to the nucleotide sequence encoding the linker sequence KKVELFPN. In the PCR process an upstream ClaI site and a downstream XhoI site were introduced by the primers allowing for ligation into the vector pGEX-4T-3 containing the Fiber A7 EGF gene construct to obtain Fiber A7 ZIgG/ZIgA [SEQ. ID. NO.: 18].

A further gene construct was also assembled to encode a fiber containing two linked ZIgG domains. This gene codes for the fiber A7 ZIgG /ZIgG [SEQ. ID. NO.: 19] and was assembled exactly as described for A7 ZIgG /ZIgA with the exception that the gene for ZIgG was used instead of ZIgA in the PCR reaction.

For a schematic representation of the constructions and construction pathways see Figs 1 and 2.

EXAMPLE 5: Binding studies

The genes encoding recombinant fibers were cloned into the vectors pCDNA (which targets proteins for expression in the cytosol) and pSecTag (which targets proteins for expression as secreted products), both from Invitrogen

- 54 -

BV, Groningen, The Netherlands, and transfected into COS7 cells using Lipofectamin (Life Technologies Inc, Gaithersburg, MD, USA) as described by the manufacturers.

5

Expression and cellular localization of recombinant fibers were evaluated by immunostaining using the following primary reagents:

10 Mouse monoclonal antibody 4D2.5 (anti-Ad5 fiber) (kindly provided by Dr Geoffrey Engler, University of Birmingham, Alabama, USA) (Shin Hong et al., Virology 185: 758-767, (1991)).

15 Mouse monoclonal antibody 2A6.36 (anti-trimerised Ad5 fiber) (kindly provided by Dr Geoffrey Engler, University of Birmingham, Alabama, USA) (Shin Hong et al., Supra).

20 Mouse monoclonal antibody against Epidermal Growth Factor (EGF) = a-EGF (Cambio, Cambridge, UK, Cat no CA 954).

Biotinylated mouse monoclonal anti idiotypic antibody directed against monoclonal antibody C242 = a-Id C242 (Lindholm et al., unpublished results).

25 Biotinylated mouse monoclonal anti idiotypic antibody directed against monoclonal antibody G250 = a-Id C250 (Kindly supplied by Reinder Bolhuis, Daniel Den Hoed Cancer Center, Rotterdam, The Netherlands).

30 Human polyclonal IgG and IgA for evaluation of Affibody activity = HIgG (Sigma-Aldrich Fine Chemicals, Cat no I4506) and HIgA (Sigma-Aldrich Fine Chemicals, Cat no I1010).

Secondary reagents were:

35 For identification of mouse antibodies: FITC labelled F(ab)2 rabbit anti mouse immunoglobulin (DAKO A/S, Glostrup, Denmark, Cat no F0313) = aMIg.

For identification of human IgG: FITC labeled F(ab)2

- 55 -

rabbit anti human IgG (DAKO A/S, Glostrup, Denmark, Cat no F0315) = aHIgG.

For identification of human IgA: FITC labeled F(ab)₂ rabbit anti human IgA (DAKO A/S, Glostrup, Denmark, Cat no F0316) = aHIgA.

For identification of biotinylated antibodies: FITC labeled Streptavidin (DAKO A/S, Glostrup, Denmark, Cat no F0422).

10 Briefly, cells were centrifuged onto microscope slides in a Shandon Cytospin2 cytocentrifuge and air-dried over night at room temperature. The preparations were fixed in 3% paraformaldehyde in phosphate buffered saline, pH 7,4 (PBS), permeabilized with 0,1% Triton-X100 in PBS.

15 After washing in PBS, preparations were incubated with primary reagents for 30 minutes at 37°C in a humid chamber, washed again in PBS and incubated with secondary reagent for 30 minutes at 37°C in a humid chamber. After washing in PBS, preparations were

20 mounted in PBS with 50% glycerol and viewed in a Zeiss Axophot microscope equipped with appropriate light source and filters for FITC.

The results are shown below in Tables 2 and 3. It is

25 obvious that all of the different ligands show appropriate binding when the corresponding fibers were expressed as secreted products. However, only the affibodies show the expected correct binding when the fibers were expressed in the cytosol and subsequently

30 transported to the nucleus. Therefore, not all ligands can fold correctly in the cytosol and nucleus. It is interesting that the ligands which can withstand the milieu in the cytosol and nucleus (affibodies) are small

35 α -helical structures not depending on S-S bridges for their conformation whereas the ligands which were shown not to be properly expressed in the cytosol/nucleus all have a conformation that is dependent on the formation

- 56 -

of S-S bridges which are formed very poorly or not at all in the cytosol.

5 **Table 2: Expression and functional ligand binding in recombinant adenovirus fibers targeted for secretion in COS cells**

Detecting reagent								
10	Fiber	4D2+	2A6+	a-EGF+	a-MIg	a-Id+	HlgG+	HlgA+
		aMIg	aMIg	aMIg		aMIg	aHlgG	aHlgA
	WT	+	+	ND	ND	ND	ND	ND
	A7 EGF	+	+	+	ND	ND	ND	ND
15	A7 G250	+	+	ND	+	+	ND	ND
	A7 C242	+	+	ND	+	+	ND	ND
	A1 ZIgG	+	ND	ND	ND	ND	+	-
	A7 ZIgG	+	ND	ND	ND	ND	+	-
	A7 ZIgG/ZIgG	+	ND	ND	ND	ND	+	-
20	A7 ZIgA	+	ND	ND	ND	ND	-	+
	A7 ZIgG/ZIgA	+	ND	ND	ND	ND	+	+

- 57 -

Table 3: Expression and functionality of ligand binding in the nucleus of COS cells in native and selected recombinant fibers after targeting for expression in the cytosol

5

Detecting reagent

	Fiber	4D2+	2A6+	a-EGF+	a-MIg	a-Id+	HigG+	HigA+
		aMIg	aMIg	aMIg		aMIga	HIGG	aHIGa
10	WT	+	+	ND	ND	ND	ND	ND
	A7 EGF	+	+	-	ND	ND	ND	ND
	A7 G250	+	+	ND	+	-	ND	ND
	A7 C242	+	+	ND	+	-	ND	ND
15	A1 ZIgG	+	ND	ND	ND	ND	+	-
	A7 ZIgG	+	+	ND	ND	ND	+	-
	A7 ZIgG /ZIgG	+	ND	ND	ND	ND	+	-
	A7 ZIgA	+	+	ND	ND	ND	-	+
	A7 ZIgG/ZIgA	+	ND	ND	ND	ND	+	+
20	A7 ZIgG Xa Knob	+	ND	ND	ND	ND	+	-

The results have obvious implications for the construction of those re-targeted virus for human gene therapy where the viral structural components containing the new cell binding ligand are synthesized in the mammalian cell cytosol, i.e. adenovirus. Below are two enabling examples to show how such re-targeted adenovirus can be constructed.

30

EXAMPLE 6:Grafting of CDR loops

35

Certain single chain constructs of monoclonal antibodies retain their binding specificity even in the mammalian cell cytosol (Cattaneo et al., TIBTECH 17: 115-121, (1999)). This is a function of the so called frame work regions of the antibody variable regions. It is known

- 58 -

that the antigen binding CDR loops can be transferred from one antibody to another by recombinant DNA technology thereby creating antibodies with frame work properties from one antibody and binding properties from another (for methodology see Emery et al., Strategies for Humanizing Antibodies, in Antibody Engineering, Carl A.K. Borrebaeck, (ed.), Oxford University Press 1995, page 159).

Such a loop-grafted single chain antibody, based on a variable domain framework capable of folding in the cytosol and subsequent transport to the cell nucleus thus created can subsequently be supplied with appropriate cloning sites at the gene level, ligated into fiber A7 RGD encoding genes and rescued into the adenovirus genome as described below (Example 7).

EXAMPLE 7:

Rescuing of recombinant fibers into the adenovirus genome

The wild type fiber as encoded in the Ad genome was substituted for recombinant fibers by the following method developed within the project. In the method the wild type Ad5 genome in the plasmid pTG3602 (Emery et al., Supra) was used as receptor for genes encoding the recombinant fibers. This plasmid contains the entire wild-type Ad5 genome joined to the plasmid backbone by Pac1 linkers. The entire genome can be recovered as a linear DNA fragment after cleavage with Pac1 since Pac1 sites are absent from the Ad genome. The resulting linear Ad DNA can then be transfected to susceptible cells to yield virus (Chartier et al., Supra). From this plasmid the Ad genome can also be cleaved as two fragments, one of 27 kb and one of 9 kb, using the enzymes Pac1 and Spe1. The 9 kb fragment has been cloned into pBluescript. From the 9 kb fragment, which

- 59 -

contains the fiber, a 3 kb SacI-KpnI fragment containing the fiber gene was further subcloned. The fiber gene was deleted between the NdeI site in the tail-portion of the fiber and the MunI site which is situated just down-stream of the fiber-gene and an adapter containing an XhoI site and the down-stream sequence was introduced between the NdeI and MunI site to obtain a fiber shuttle vector. Several recombinant fibers have now been ligated between NdeI and XhoI of this shuttle vector and thereafter rescued into the 9 kb fragment mentioned above by homologous recombination in *E. coli* (Chartier et al., Supra). This means that all normal elements regulating fiber-expression have been left intact.

Finally, recombinant 9 kb fragments were separately joined to the 27 kb fragment by cosmid cloning to re-create the complete Ad genome.

The 27 kb fragment may also be derived from another adenovirus genome, such as the Ad-YFG described by He et al (He T-C, Zhou S, DaCosta LT, Yu J, Kinzler KW and Vogelstein B: A simplified system for generating recombinant adenoviruses. Proc Natl Acad Sci, USA, 95:2509-2514, 1998). If the 27 kb fragment is derived from pTG3602 the resulting genome will be WT E1+ whereas the 27 kb fragment from Ad-YFG will render E1 deleted viruses needing e.g. low passage 293 for replication.

The recombinant Ad genome resulting from above manipulations can finally be obtained as a linear fragment by cleaving with PacI and used to transfect permissive cell lines which then yield virus plaques if the genome is functional. For these transfections the FuGENE 6 transfection agent (Roche) was used.

Various recombinant fibers were rescued into the Ad genome and subsequently transfected into permissive

- 60 -

cells. Results are shown below in table 4. Of the fibers accounted for, only those containing the WT knob and the affibody A7 ZIgG/IgG were capable of rendering functional virions. The results are in complete
 5 concordance with those shown in tables 2 and 3.

Table 4: Rescuing of recombinant fibers into virions

10	<i>Fiber</i>	<i>Transfected cells</i>	<i>Occurrence of virus plaques</i>
	A7 WT Knob	293	Yes
	A7 EGF	A549	No
	A7 C242	Colo 205	No
	A7 G250	A75	No
15	A7 ZIgG/ZIgG	293/Fc*	Yes

(*) 293/Fc cells are 293 cells stably transfected with Fc3(1) from human IgG expressed as a membrane protein. To create 293/Fc, cloned Fc3(1) from human IgG which
 20 reacts with the Z-domain (Jendeberg, L., Nilsson, P., Larsson, A., Nilsson, B., Uhlén, M. and Nygren, P.-Å (1997) "Engineering of Fc1 and Fc3 from human immunoglobulin G to analyse subclass specificity for staphylococcal protein A" J. Imm. Methods 201, 25-34.)
 25 was ligated into the vector pDisplay (Invitrogen) in frame with the PDGFR transmembrane domain sequence present in this vector. The coding sequence containing the Fc sequence fused to the PDGFR transmembrane domain sequence was then cleaved from the vector using the
 30 restriction enzymes SfiI and NotI and ligated into the vector pSecTag (which carries the Zeocin resistance gene) cleaved with the same enzymes. The reconstituted vector was transfected into low passage 293 cells using FuGENE and the cells were placed under selection
 35 pressure using Zeocin to select for stably transformed cells. Clones were isolated and tested for membrane expression of membrane-bound Fc3(1) using FITC labeled

- 61 -

staphylococcal protein A (Sigma). One 293 clone which homogenously expresses membrane bound Fc was used for transfection of the A7 ZIgG/ZIgG containing genome to produce virions.

5

EXAMPLE 8:

Assembly of gene constructs encoding recombinant adenovirus fibers with an affibody and a cleavable wild type knob

10

Cleavable fibers containing both a non-native polypeptide comprising an external ligand, here an affibody, and a C-terminally placed wild type knob were constructed with an activated factor X site situated between the cell binding structures so that the knob can be cleaved off to expose the affibody. This permits virus production in 293 cells with subsequent infection of new target cells as defined by the affibody after proteolytic removal of the knob.

20

To construct a gene for a cleavable fiber for ligation of different ligands the WT fiber was subjected to PCR with an upstream primer (437) introducing ClaI, MunI and the Factor Xa recognition site before the first seven triplets of repeat 22 and the downstream primer 149 which primes at the end of the fiber knob. After cloning and restriction with ClaI and XhoI the DNA fragment was cloned into the A7 fiber construct mentioned earlier. The resulting fiber gene contains from the N-terminus the sequence for the tail, the first seven shaft repeats, the NRP trimerisation signal, the linker from staphylococcal protein A, a ClaI site, a MunI site, the Xa cleavage site, repeat 22 and the wild type knob. For ligation into this fiber gene the gene for the affibody ZIgG was supplied with an upstream ClaI site and a downstream MunI site by PCR using primers 503 and 505 and ligated into the aforementioned "Xa fiber"

- 62 -

gene. The resulting fiber gene contains from the N-terminus the sequence for the tail, the first seven shaft repeats, the NRP trimerisation signal, the linker from staphylococcal protein A, a ClaI site, the new
5 linker, a MunI site, the Xa cleavage site, repeat 22 and the wild type knob.

The fiber gene was rescued into the adenovirus genome (see Example 7). The recombinant genome was transfected
10 into 293 cells and virions were produced and purified on CsCl gradients. Purified virions were cleaved with activated Factor X (Xa) (Sigma) in 50mM Tris-Cl pH 8,0; 100 mM NaCl; 5mM CaCl₂ for 48 hours at room temperature. 1 U of Xa was used for 5µL of virus suspension.

15 Assay for binding of the ZIgG affibody on the virions was performed essentially as follows. Virus suspensions were blotted onto PVDF membranes (BioRad). After blocking with 3% gelatin in Tris buffered saline, pH 7,4
20 (TBS) the membranes were incubated with FC 10µg/mL in 1% gelatin in TBS for 90 minutes at room temperature, washed, incubated with anti-human IgG-HRP (Dakopatts) in TBS/1% gelatin for 90 minutes at room temperature, washed and developed in 4-Chloro Naphtol reagent
25 (BioRad). The results of staining of WT and recombinant viruses are shown in Figure 3 along with different controls. It is obvious that the recombinant virus binds Fc3(1) which binds to WT Z whereas Fc3 which does not bind to WT Z (Jendeberg, L., Nilsson, P., Larsson, A.,
30 Nilsson, B., Uhlén, M. and Nygren, P.-Å. (1997) "Engineering of Fc1 and Fc3 from human immunoglobulin G to analyse subclass specificity for staphylococcal protein A" J. Imm. Methods 201, 25-34) fails to bind to the recombinant virus. WT virus does not bind either Fc
35 preparation.

Example 9:Assay of non-native polypeptides for functional conformation and binding specificity

5

The candidate polypeptide is expressed as part of a viral component protein by means of insertion of the corresponding coding sequence into a suitable construct. The construct, which comprises an entire viral genome in which one or more components have been replaced by the corresponding recombinant component gene, is expressed in a cell line suitable for propagation of the recombinant virus coded for by the recombinant viral genome. There are two main possibilities depending on whether the WT cell-binding function is retained in the recombinant genome or not.

15

If the WT cell-binding function is retained virus can be produced in normal producer lines such as 293 for adenovirus. The ability of the non-native polypeptide to be expressed in a conformation that allows it to bind to an extracellular ligand and to be part of a functional virion is assayed by binding studies on virions. This approach is exemplified by the strategy employed in Example 8 above where the virus fiber contains both the WT knob and a non-native binding ligand and where the knob can be removed by proteolytic cleavage. The binding function of the non-native polypeptide was demonstrated by standard solid phase immunotechnology. Another strategy to meet the same ends would be to produce virions with WT fibers as well as recombinant fibers, the latter containing the non-native polypeptide as replacement for the native knob.

20

25

30

If the WT cell-binding function has been deleted from the viral genome the virions will depend on the non-native polypeptide for cell binding. Therefore, the

35

- 64 -

cell line must be able to permit entry of the recombinant virus and, if necessary, to supply genetic information that may be missing from the recombinant virus, e.g. the sequence for E1. If the recombinant genome is E1 deleted, one solution is to use an E1 transfected cell line such as 293 and to stably transfect these with a receptor capable of binding the non-native polypeptide coded for by the recombinant viral genome. The ability of the non-native polypeptide to be expressed in a conformation that allows it to bind to the corresponding receptor structure and form part of a functional virion is assayed by screening for plaque formation in the cells following transfection of the viral genome. This strategy was employed in Example 7 above where 293 cells stably expressing membrane bound Fc from human IgG was used to produce virions where the WT binding function was replaced with the affibody ZIgG. The exact binding properties of the non-native polypeptide can further be determined in studies where the virus is allowed to compete for binding with peptides structurally related to the non-native polypeptide responsible for cell-binding.

EXAMPLE 10:

Anti β -galactosidase single chain Fv fragment

In a continuation of the experiments described in Example 3, a single chain variable chain fragment (VK, linker, VH) reactive with β -galactosidase and capable of being expressed in the cytoplasm (Martineau P and Betton J-M: J. Mol. Biol. 292, 921-929 (1999)) was cloned into the aforementioned A7 fiber constructs. This single chain fragment is [SEQ. ID. NO.: 47]. To permit cloning into the aforementioned A7 fiber constructs, the single chain fragment was cut with NcoI/EcoRI from the plasmid pPM163R4 (Martineau P and Betton J-M *supra*) and ligated into a plasmid containing the aforementioned Fiber R7

- 65 -

EGF supplied with a ClaI-NcoI-EcoRI-XhoI adapter allowing for in-frame ligation of the single chain fragment.

5 In a continuation of the experiments described in Example 5, expression and cellular localization of recombinant fibers containing the single chain variable chain fragment (VK, linker, VH) reactive with β -galactosidase were evaluated by immunostaining using
10 the following additional primary reagent:
 β -galactosidase-biotin (SIGMA, G 5025).

In addition to the results shown in Table 2, results
15 results obtained in equivalent experiments for the Anti β -galactosidase single chain Fv fragment fiber construct are shown Table 5:

20 **Table 5: Expression and functional ligand binding in recombinant adenovirus fibers targeted for secretion in COS cells**

Fiber	4D2+ aMIg	2A6+ aMIg	β -galactosidase + StrAv-FITC
R7 a- β -galactosidase	+	+	+

25

In addition to the results shown in Table 3, results
obtained in equivalent experiments for the Anti
 β -galactosidase single chain Fv fragment fiber construct
are shown Table 6:

30

Table 6: Expression and functionality of ligand binding in the nucleus of COS cells in native and selected recombinant fibers after targeting for expression in the cytosol

5

Fiber	4D2+ aMIg	2A6+ aMIg	β -galactosidase + StrAv-FITC
R7 a- β -galactosidase	+	+	+

10

EXAMPLE 11:

Phenotypic analysis of fiber proteins.

15 The following is a further example of modified viruses of the present invention wherein complex polypeptide ligands have been incorporated into a modified
adenoviral fibre protein. The example demonstrates the importance for generation of a re-targeted viable and
20 functional Ad vector of two features: (i) the fibre structure modifications should still allow for efficient attachment and cellular entry of the virus, and (ii) ligands inserted into the fibre should be capable of correct folding in the mammalian cell cytoplasm.
25 Solubility is conveniently used to assess correct folding which is typically linked to an absence of disulphide bonds. Preferably the non-native polypeptides described herein will have no more than 2 disulphide bonds, typically no more than 1 and most
30 preferably no disulphide bonds.

Materials and Methods

Cells. HEK-293 cells were obtained from Microbix Inc.
35 (Toronto, Ontario, Canada). Cos7, A431 and Colo205 cells were purchased from the American Type Culture Collection

(ATCC; Manassas, Virginia) and A75 and G43 cells were obtained from Reinder Bolhuis (Daniel Den Hoed Cancer Center, Rotterdam, The Netherlands). All cells were maintained at 37°C and 5% CO₂ in Iscove's medium (Gibco BRL), supplemented with 10% fetal bovine serum (Sigma-Aldrich) and 50 mg/ml Gentamicin (Gibco BRL). Spodoptera frugiperda (Sf9) cells (ATCC) were cultured at 28°C in TC 100 medium (Gibco BRL) with the same supplements as above.

10

Antibodies. Monoclonal antibody (mAb) against epidermal growth factor (EGF) was purchased from Cambio Ltd. (Cambridge, UK). Antibodies directed against the V α and V β domains of single chain T-cell receptor (scTCR), and anti-idiotypic antibodies directed against mAb G250 (NUH31 and NUH84) were kindly supplied by Reinder Bolhuis. MAb against fiber tail (4D2.5), and mAb against fiber trimer (2A6.36) were obtained from Jeff Engler (University of Alabama at Birmingham, AL). CAR-blocking, fiber knob directed mAb 1D6.14 was supplied by Buck Rogers (UAB at Birmingham, AL). The monoclonal antibody RL2, which is specific for O-linked GlcNAc residues, was obtained from Larry Gerace via Jeff Engler. Biotinylated anti-idiotypic antibodies directed against mAb C242 (Id1, Id13 and Id20) were produced from the original hybridomas. Horse radish peroxidase (HRP)-labeled streptavidin, fluorescein isothiocyanate (FITC)-labeled rabbit anti-mouse immunoglobulin G and streptavidin-FITC were purchased from DAKO (Glostrup, DK).

30

Generation of recombinant knobless fibers and nomenclature. Recombinant fiber genes were constructed using methods based on ligation, PCR, and splicing by overlap extension (SOE). Gene sequences generated by PCR were sequenced before subcloning. The gene encoding the Ad5 WT fiber was obtained from pAB26 (Microbix, Toronto, Canada) by PCR using the forward primer (SEQ ID NO: 48)

35

- 68 -

5'-CTC GGA TCC GAT GAA GCG CGC AAG ACC GTC TGA A-3' and reverse primer (SEQ ID NO: 49) 5'-TTC CTC GAG TTA TTC TTG GGC AAT GTA TGA-3' introducing an upstream Bam H I and a downstream Xho I site, respectively. In

5 recombinant fibers, the knob domain was deleted and replaced by a 36 amino acid (aa) extrinsic trimerisation motif derived from the neck region peptide (NRP) of human lung surfactant protein D. The NRP sequence (PDVASLRQQVAELQGQVQHLQAAFSQQYKKVELFPNG) (SEQ ID NO:2),

10 followed by a linker sequence from Staphylococcus protein A (Staph-A linker: AKKLNDQAQPKSD), was ligated to the C-terminal end of fiber shaft of different lengths, 1, 7 or 22 repeats.

15 The resulting constructs were named R1, R7 and R22, respectively. Re-targeting ligands were added to the C-terminal end of the Staph-A linker. For convenient cloning of various ligands, Cla I and Xho I sites were introduced after the linker sequence. All ligands

20 mentioned below were provided with these restriction sites, and the name of the ligand was indicated after the number of shaft repeats in the fiber name. E.g., R1-RGD, R7-EGF, etc (Fig. 4). R7-knob referred to a truncated fiber shaft (repeats 1 to 7), carrying NRP,

25 the Staph-A linker and the natural knob domain, including the last shaft repeat and the shaft-knob junction (Fig. 4). For two R7-knob fiber constructs, an extra linker was inserted into the Nhe I site located on the N-terminal side of NRP. These linkers were derived

30 from the Ad5 WT fiber shaft repeat 17 (L17: TTTACAGCTTCAAACAATTCCAAAAAGCTTGAG) and fiber shaft repeat 22 (L22 GGAAACAAAAATAATGATAAGCTAACTTTGTGTGACC) were named as R7-L17-knob and R7-L22-knob, respectively. The schematic representation of the fibers is shown in

35 Fig 4.

Targeting ligands. Polypeptide ligands. Double stranded

- 69 -

DNA fragments containing the RGD or ACDCRGDCFCG (abbreviated RGD4C) motifs were synthesized as complementary oligonucleotides with single-stranded terminal adapters, and annealed together. The fragment containing the shaft repeat 22, the shaft-knob junction and the entire knob was obtained by PCR from the WT Ad5 fiber gene. The DNA sequence for human EGF was synthesized, cloned and verified by sequencing. The cloned single chain T-cell receptor (which contains disulfide bonds) with specificity towards Magel/HLA A1 was obtained from Reinder Bolhuis. The scTCR used had the format V α -Linker-V β C β .

Monoclonal antibodies. Two mAb single chain fragments (scFv) were used. The first scFv consisted of the variable kappa chain (or VK), a spacing linker, the variable heavy chain (or VH), the joining sequence (JS) and the constant heavy chain domain 2 (or CH2) of mAb G250. MAb G250 has been shown to have a high specificity towards a surface antigen of human renal carcinoma cells.

Affibodies. The second scFv contained the same basic structure VK-linker-JS-VH, and derived from mAb C242, which reacts with colorectal and pancreatic carcinomas. However, to obtain a better selectivity for the designed target cells, a library of C242 scFv-derived affibodies was generated by SOE, using cDNA from the mAb-producing hybridoma cells as the original template. A randomized 18 amino acid peptide ligand (SNN)₁₈ was inserted between the VK and VH domains, where S was dA, dG or dC and N was dA, dG, dC or dT. Affibodies were selected by phage display on colorectal carcinoma cells Colo205, and the sequence PPDFVPPAASFDPHSPRG was identified for the peptide ligand. The C242 scFv VK-(PPDFVPPAASFDPHSPRG)-VH was then selected for further work based on its antigen binding ability.

- 70 -

Cellular expression and localization of fiber and ligand reactivity. The genes encoding recombinant fibers were cloned into the vectors pCDNA3.1 and pSecTag2 (Invitrogen BV, Groningen, Germany) for intracellular expression and extracellular release, respectively. Vectors were transfected into Cos7 cells using Lipofectamin (Life Technologies Inc., Gaithersburg, MD, USA) following the manufacturers' instructions. Expression, nuclear transport and functional expression of the fibers with the new cell binding ligands were evaluated by immunostaining as follows. Cells were centrifuged onto microscope slides in a Shandon Cytospin2 cytocentrifuge and air-dried over night at room temperature. The preparations were fixed in 3% paraformaldehyde in PBS and permeabilized with 0.1% Triton-X100 in PBS. After washing in PBS, preparations were incubated with primary antibodies (anti-fiber mAb, or anti-ligand mAb) for 30 min at 37°C in a humid chamber, washed again in PBS and incubated with secondary antibodies labeled with FITC for 30 min at 37°C in a humid chamber. After washing in PBS, preparations were mounted in PBS with 50% glycerol and viewed in a Zeiss Axioskop microscope equipped with appropriate light source and filters for FITC.

Phenotypic analysis of fiber proteins. Recombinant fiber proteins were expressed in insect cells infected with recombinant baculoviruses, and analysed according to four criteria: (i) solubility, (ii) trimerization, (iii) glycosylation and (iv) assembly with recombinant penton base in vivo to form penton capsomers.

(i) For immunological quantification of soluble versus insoluble recombinant fiber fractions, Sf9 cells were lysed in hypotonic buffer (10 mM Tris-HCl buffer, pH 7.5) at 0°C, and the cell lysates were adjusted to isotonic conditions (150 mM NaCl in 10 mM Tris-HCl, pH

- 71 -

7.5) and subjected to centrifugation at 15,000 x g for 10 min. Supernatants and pellets were then analyzed by conventional SDS-PAGE and immunoblotting, using anti-tail 4D2.5 mAb as primary antibody, and
5 [35S]SRL-labeled anti-mouse IgG secondary antibody (Amersham Pharmacia Biotech; 100 µCi/ml; 5 µCi per blot). (ii) Oligomerization status of fiber was assayed by means of non-denaturing SDS-PAGE (referred to as NDS-PAGE) and conventional, denaturing SDS-PAGE.
10 NDS-PAGE differed from SDS-PAGE in that the samples were not denatured by boiling in SDS sample buffer prior to electrophoresis. (iii) Glycosylation of recombinant fibers was assessed both by immunoreaction on blots using the monoclonal antibody RL2 and chemical detection
15 using the DIG Glycan Detection Kit (Roche). (iv) Assembly of fiber with penton base was assayed by co-infecting the same Sf9 cells with two recombinant AcNPV, one expressing the penton base, the other expressing the fiber protein.
20
The presence of penton capsomer was detected in cell lysates after 40 h post-infection, and analysed by PAGE in native conditions, at low voltage overnight with cooling. Immunological quantification of native penton,
25 penton base and fiber proteins was performed as above, using the corresponding primary antibody (anti-penton base or anti-fiber), followed by [35S]SRL-labeled anti-mouse or anti-rabbit whole IgG secondary antibody. Blots were exposed to radiographic films (Hyperfilm
30 beta-max, Amersham Pharmacia Biotech), and autoradiograms were scanned at 610 nm, using an automatic densitometer (REP-EDC, Helena Laboratories, Beaumont, TX). Alternatively, protein bands were excised from blots and radioactivity measured in a scintillation
35 counter (Beckman LS-6500).

Rescue of recombinant modified fibers into virions.

- 72 -

Recombinant fibers were rescued into the Ad5 genome as described above. Briefly, a 9 kb-fragment from the Ad5 genome, spanning map units (mu) 75.4 to 100 and containing the WT fiber gene, was generated by Spe I and Pac I digestion of pTG3602. The plasmid pTG3602 contained the entire WT Ad5 genome bounded by two Pac I sites. The 75.4-100 mu-fragment was then cloned into pBluescript II SK(-) (Stratagene), after addition of a Pac I site in its MCS, generating the plasmid pGAG9. From pGAG9, a Sac I-Kpn I 3kb-fragment was then subcloned into pBluescript II SK(-). A large deletion downstream of the Nde I site (located within the tail domain of Ad5 fiber) was created in the 3kb-fragment, and the deleted sequence replaced by a Xho I site-containing linker. This generated the plasmid pGAG3, which was the receiving plasmid for all our fiber gene constructs, inserted between Nde I and Xho I sites. pGAG3-inserted recombinant fibers were re-introduced into pGAG9 digested with Nhe I, using homologous recombination in E. coli BJ5183. The resulting recombinant pGAG9 was then excised from the vector with Spe I and Pac I, and joined to the isolated 27 kb-fragment (0-75.5 mu) representing the left-hand segment of the Ad5 genome by Cosmid cloning (SuperCos 1 Cosmid Vector Kit and Gigapack III Gold Packaging Extract, Stratagene). The presence of the correct recombinant fiber in the cosmid clones were verified by PCR and restriction analysis using Spe I and Hind III.

For virus production, recombinant cosmid genomes were isolated after restriction with Pac I, and transfected into cells expressing the receptors corresponding to the fiber-inserted ligand. In standard transfection reactions, 2 μ g DNA and 3 μ l FuGENE (Roche) were used per 35-mm well, according to the manufacturer's protocol. Ad5 genomes with fiber gene containing either RGD motifs or the WT fiber knob as ligands were

- 73 -

transfected into 293 cells. For the other liganded fibers, Ad5 genomes with the EGF ligand were transfected into A549 cells, scFv-C242 into Colo 205 cells, scFv-G250 into A75 cells and scTCR specific for
5 Magel/HLA A1 into G43 cells. At least three different transfections were simultaneously performed, each one using a 6-well plate. The occurrence of plaques was determined by microscopic observation of the transfected cell cultures. Verification of recombinant fiber
10 sequence was made by PCR with specific primers for each fiber construct.

Characterization of recombinant fiber and Ad5 virions.

15 Cellular expression. To evaluate the level of fiber expression, 293 cells were infected with 10 pfu/cell of WT, R7-knob and R7-RGD virus. Cells were harvested and freeze-thawed four times analyzed by SDS-PAGE and western blotting. The blots were reacted with 4D2.5 and
20 revealed with HRP-labeled anti-mouse IgG (DAKO).

Fiber content of Ad5 virions. The fiber copy number of virions was determined, after CsCl purification of Ad5 virions, by SDS-PAGE and western blot analysis as above.
25 The virus loads in acrylamide gels were normalized for equal amounts of infectious particles, determined by virus titration on 293 cells (expressed as plaque forming units/ml; PFU/fml), and for equal amounts of physical particles (PP), determined by protein assay
30 (BioRad). The number of PP was determined by SDS-PAGE analysis and Coomassie blue staining of Ad5 recombinants, co-electrophoresed with a range of standard bovine serum albumin (BSA) samples (2x crystallized, BioRad). Protein content of hexon bands
35 was evaluated by comparison with BSA standard bands, by scanning in an automatic densitometer (REP-EDC, Helena Laboratories, Beaumont, TX). The number of Ad5 PP in

- 74 -

samples was calculated from the mass of 2.91×10^{-16} g per single virion, i.e. 2.91 mg per 10^{+13} virions. The infectivity index represented the ratio of infectious to physical particles.

5

Growth rate of recombinant Ad5. Growth-rate was measured by the production of plaques in 293 cells. In standard assays, virus in PBS was adsorbed to cell monolayers (4×10^4 cells per sample) at 37°C for 1 h. The cells were rinsed once and further incubated in Iscove's medium supplemented with 10 % FCS and 50 mg/ml gentamycin at 37°C. Cells were harvested at 24, 48 and 72 h after infection (pi), centrifuged, dissolved in 0.2 ml PBS and freeze-thawed four times. The supernatants were titered on 293 cells, and the titer expressed as plaque forming units per ml. The total Ad protein content was measured by the IDEIA™ Adenovirus kit (DAKO).

10

15

Gene transduction efficiency. Monolayers of 293 cells in 24-well plates were infected as described above with 10 pfu/cell of the recombinant viruses. Cells were harvested at 24 h pi, washed with ice-cold PBS followed by fixation in 0.5 % glutaraldehyde for 15 min. After three washes in PBS, the cells were analysed for transgene GFP expression, using the FL1 emission channel in a FACScan cytometer (Becton-Dickinson, San Jose, CA).

20

25

Assay for fiber knob in viral capsids. The presence or absence of accessible knob in virions was assessed by ELISA. Purified virions were diluted with 50 mM carbonate-bicarbonate buffer pH 9.6, to a final concentration of 5×10^5 PFU/ml. Aliquots of 100 ml were adsorbed onto ELISA plates overnight at 4°C. Material adsorbed in wells was fixed with 0.5% glutaraldehyde, after which the wells were washed with PBS containing 0.1% Tween-20 (washing buffer), and blocked for 2 h with 200 ml PBS containing 1% BSA. Wells were then washed

30

35

- 75 -

three times in washing buffer and incubated for 1 h at room temperature with 100 ml of biotinylated anti-knob mAb 1D6.14 at a concentration of 1 mg/ml. Bound antibody was detected using HRP-streptavidin (DAKO) at a dilution of 1:2,000, for 1 h at room temperature. Colour development was obtained with TMB substrate (CanAg Diagnostics, Göteborg, Sweden), and stopped with 0.12 M HCl for 10 min. Plates were read in a microtiter plate reader set at 450 nm.

10

Rationale for the construction of recombinant fibers.

Knobless fibers with the extrinsic trimerisation motif (NRP) from human lung surfactant protein D were constructed with different numbers of shaft repeats and different ligand structures. The three constitutive elements of these fiber constructs were considered from a structural and functional point of view: (i) the fiber scaffold, (ii) the flexible linker and (iii) the cell ligands.

20

Fiber scaffold. The fibers were named according to the number of shaft repeats and the ligand present. As an example, R7-EGF contained the fiber tail, the N-terminal 7 shaft repeats, the NRP motif and the C-terminal EGF peptide as the cellular ligand. R1-RGD fiber had only one shaft repeat (the first one) and RGD as the ligand, etc. Of three possible shaft lengths, short (1 repeat), medium (7 repeats) and long (22 repeats) repeats, the intermediate size fiber with 7 repeats (R7) was chosen as the building scaffold for further constructions and studies with different ligands. For reasons discussed below, the rationale for choosing R7 was based on its high solubility as recombinant protein, the maintenance of most of the fiber biological functions, and on the yields of Ad5/FibR7 virus progeny. The study on the comparative advantages of fiber shafts of different

35

- 76 -

lengths was performed with R1-RGD, R7-RGD and R22-RGD fibers, three constructs carrying the same cellular ligand RGD.

5 **Linker.** To evaluate the possible advantage of an additional linker in the fiber shaft domain, two R7-NRP-knob fibers were constructed with an extra peptide linker inserted between the shaft domain and the NRP motif. Two virus-derived linkers were thus tested,
10 the Ad5 shaft repeat 17 (L17) and the Ad5 shaft repeat 22 (L22). The resulting recombinant fibers were named R7-L17-knob and R7-L22-knob, respectively.

15 **Cellular ligands.** Several ligands were tested and compared for functionality. These ligands were designed to re-targeted Ad5 vectors to cell surface molecules of broad distribution, like integrins or HLA molecules, or to less ubiquitous molecules, like malignant cell
20 specific determinants. They varied in size and complexity from a simple tripeptide motif, like RGD, to more elaborated structures, like scTR or scFv, which consist of several polypeptide domains with requirement for proper folding.

25 **Expression of fibers and ligands in mammalian cells.** Recombinant fibers were first transiently expressed in mammalian cells, using plasmid-transfected Cos7 cells. The recombinant fibers were expressed from two different
30 vectors, one designed for intracellular expression of recombinant proteins, the other for their extracellular release via the secretory pathway. Cells were assayed for the level of fiber expression, cellular localization and functionality of their ligands. All the different
35 fibers tested exhibited apparent appropriate ligand binding when recovered as secreted proteins. For fibers synthesized from the intracellular expression vector, nuclear localization was observed in all cases, except

- 77 -

for R7-L22-knob. This implied that the cytoplasmic transit and the traverse of the nuclear pore occurred as efficiently for the modified fibers as for WT Ad5 fiber.

5 However, none of the extrinsic ligands fused to fibers showed the expected binding activity. Even though the intracellularly expressed R7-scTCR fiber could be detected within the nucleus, and was stained with an anti-Va mAb recognizing an epitope independent of the Va
10 domain conformation, no staining could be detected with another, conformation-dependent mAb, directed against the Vb domain. In order to test the possible detrimental effect of the fiber domain on the reactivity of intracellular ligands, our ligands were also expressed
15 as separate constructs from the vector pcDNA. There was no detectable binding of the nonfused intracellular ligands to their specific mAbs, indicating that the ligands behaved similarly when free or fused to the fiber.

20 There was therefore a profound difference in binding ability between fiber-ligand fusions designed for secretion and designed for cytoplasmic expression. This suggested that fiber-ligand fusion proteins designed to
25 follow the secretory pathway underwent a proper folding and were in the correct conformation. This did not imply that all intracellular fiber-ligand fusion proteins would fold incorrectly, since the cellular environment also seemed to play a major role in this process. A
30 significant difference could be detected between the cytosol of Cos7 and Sf9 cells, in terms of folding pattern and reactivity of fiber-ligand fusion proteins. This was the case for the fiber knob domain of R7-L22-knob, the EGF ligand of R7-EGF and the scFV
35 ligand of R7-G250, which showed some reactivity in Sf9 cells but not in Cos7 cells.

Phenotype of recombinant fibers expressed in insect cells. To further analyze the properties of the liganded fibers, our different constructs were expressed as recombinant proteins in the baculovirus-insect cells system, and assayed for protein solubility, trimerisation, glycosylation and formation of penton by assembly with penton base in vivo in Sf9 cells.

Protein solubility and conformation. The solubility of recombinant proteins is usually considered as a good indicator of their proper folding. We therefore tested all our recombinant fiber proteins for their total expression in insect cells, and determined the proportion recovered in the soluble fraction of the cell lysates. All the fiber recombinants were highly expressed in Sf9 cells, at levels similar to WT Ad5 fiber, although their degree of solubility varied significantly from one to another. We estimated that fibers had a WT-like solubility and thus a proper folding when the soluble fraction contained more than 50 % of the total fiber expressed. A fiber could be considered as misfolded when its soluble fraction represented less than 20 % of the total. According to these criteria, WT, R7-knob, R7-L17-knob, R7-L22-knob, R1-RGD, R7-RGD and R7-RGD4C fibers were mainly recovered in the soluble fraction (60-95 % solubility). In contrast, only 22 % of R22-RGD, and less than 15 % of R7-EGF, R7-C242 and R7-scTCR fiber was found to be soluble, confirming their incorrect folding suggested by the absence of reactivity with their respective mAbs. The least soluble fiber constructs was R7-G250, which was recovered at 95 % in the insoluble fraction (see Fig. 5 and Table 7 below).

Table 7. Phenotypic characterization of recombinant fibers expressed in baculovirus infected Sf9 cells (a)

5

Trimerization (b)					
Fiber	Glycosylation	NDS-PAGE	IF	Assembly (c) with penton base	Solubility (d) (%)
WT	+	+	+	+	60
R7-knob	+	+	+	+	72
R7-L17-knob	+	+	+	+	70
10 R7-L22-knob	+	+	+	+	75
R1-RGD	-	-	+	+	89
R7-RGD	-	+	+	+	94
R22-RGD	-	+	+	+	22
R7-RGD4C	+	+	+	+	78
15 R7-EGF	-	ND (e)	+	ND	13
R7-C242	-	ND	-	ND	11
R7-G250	-	ND	-	ND	5
R7-scTCR	-	ND	+	ND	12

20

(a) Baculovirus-infected Sf9 cells were harvested at 48 h after infection and recombinant fibers assayed for different biological functions, solubility, O-GlcNAc glycosylation, trimerisation, and penton capsomer formation.

25

(b) Trimerisation status of fibers was determined in vitro by electrophoresis of Sf9 cell lysates in SDS-gel without heat denaturation (NDS-PAGE), and in situ, by immunofluorescence staining of fixed cells, using anti-trimer mAb (2A6.36).

30

(c) The capacity of fibers to assemble with penton base to form penton capsomer was assayed by co-infection of

- 80 -

the same Sf9 cells by two different recombinant baculoviruses, one expressing Ad fiber, the other Ad penton base. Sf9 cell lysates were analysed at 48 h after co-infection for the occurrence of penton (base + fiber), by electrophoresis of native proteins in nondenaturing 6 % acrylamide gels (Karayan et al., 1994) (d) Solubility was assayed in single baculovirus-infected Sf9 cells. Cells harvested at 48 h after infection were lysed in isotonic buffer, and cell lysates divided in two aliquots. One aliquot was centrifuged at 10,000 x g for 10 min and supernatant was kept, whereas the other was analysed as whole cell lysate. Both aliquots were denatured at 100°C in SDS-sample buffer and electrophoresed in denaturing SDS-gel, and blotted. Blots were reacted with anti-fiber-tail mAb 4D2.5, and radiolabeled secondary antibody, as shown in Fig. 5. Immunoblots were quantitated by autoradiogram scanning and the results were expressed as the percentage of soluble versus total fiber content. Average of three determinations; SD was within 15 % of the reported value for the mean. (e) ND, not detectable.

Trimerisation. The ability of the different fibers to self-assemble into trimers were electrophoretically determined in vitro by NDS-PAGE analysis of cell lysates, and immunologically in situ by immunofluorescence staining of fixed cells. According to their electrophoretic patterns, R7-EGF, R7-C242, R7-G250 and R7-scTCR were incapable of forming homotrimers, whereas R7-knob, R7-L17-knob R7-L22-knob, R7-RGD, R22-RGD and R7-RGD4C trimerized at WT levels (Table 7). However, immunofluorescence staining of cells using the anti-trimer mAb 2A6.36 showed that all except R7-C242 and R7-G250 formed trimers (Table 7). The apparent discrepancy likely resided in the method of detection, or in the possibility that R7-EGF, R7-C242, R7-G250 and

- 81 -

R7-scTCR fiber trimers became unstable in vitro, or both. Whatever the reason, if fiber trimers formed with a very low efficiency within the cells, immunofluorescence staining was, in this case, more sensitive and more appropriate for trimer detection.

Glycosylation. Likewise, the glycosylation of fibers was analysed by western blotting using RL2, a mAb specific for peptide-linked O-GlcNAc residues. The fiber constructs which reacted with RL2 were, besides the WT, R7-L17-knob, R7-L22-knob, R7-knob and R7-RGD4C (Table 7).

Assembly with penton base. When Sf9 cells were infected with two different baculoviruses, one expressing the Ad fiber, the other the penton base, the two proteins are capable of interacting intracellularly to form penton capsomers. When assayed for this property, the fiber proteins R7-EGF, R7-C242, R7-G250 and R7-scTCR had lost their capacity to assemble with penton base, whereas the other eight recombinant fibers had conserved their assembly function (Table 7).

Rescue of recombinant fiber genes into the viral genome and viability of the viruses. Our recombinant fiber genes were reinserted into the Ad5 viral genome in replacement of the WT fiber gene. Each recombinant viral genome was then introduced by transfection into the corresponding cell line which express the proper receptor for the recombinant fiber. The rescue of viable recombinant viruses was assessed by plaque development. Plaques were observed for the Ad5 genomes harboring the WT fiber, and recombinant fibers R7-knob, R7-L17-knob, R1-RGD, R7-RGD and R7-RGD4C.

35

Fiber content, infectivity and growth rate of recombinant viruses. The recombinant R7-RGD Ad carrying

- 82 -

seven shaft repeats had been selected as the best choice in terms of fiber trimerisation, stability, assembly with penton base and ligand binding. However, this knobless Ad was less efficient in virus assembly and production, compared to WT virus. In this example, the characteristics of the Ad-R7-knob, which carried the knob domain, were compared to those of WT Ad and knobless Ad-R7-RGD. We first assayed for the production of fibers in 293 cells infected with each of the three Ad at different times pi. The production of fiber proteins from R7-knob Ad was comparable to that of the WT Ad, whereas the level of fiber protein from the R7-RGD Ad was significantly lower.

We then compared the fiber content per virion of the R7-knob Ad with that of the WT Ad. The two viruses exhibited similar amounts of fiber content when normalised to the number of physical particles (PP) but there were more fiber protein in the R7-knob Ad when the two virus samples were normalised to the number of infectious particles (PFU). This indicated that R7-knob Ad was less infectious than the WT. This was confirmed by the infectivity index (PFU : PP ratio), which was calculated from the infectious titer (expressed as PFU/ml, as determined by titration on 293 cells), and from the concentration of PP (determined by biochemical methods). The infectivity index of the Ad-R7-knob (1:100 to 1: 200) was 2- to 8-fold lower than that of WT Ad5 (1:25-1: 50). The infectivity index of Ad-R7-RGD (1 : 200 - 1 : 500) was in a similar range as that of Ad-R7-knob.

The growth rate of the Ad-R7-knob was compared to that of WT virus. Aliquots of cell samples were infected at the same MOI, and infectious virus progeny, determined by plaque assays on 293 cells. The growth curves for the three viruses were similar, but the production of

- 83 -

infectious Ad was inferior to the WT for both Ad-R7-knob and Ad-R7-RGD. Virus uptake was also examined for Ad-R7-knob, and estimated from its ability to transduce the GFP reporter gene into 293 cells, as compared to Ad5-GFP carrying WT fibers (Ad-WTFib). The transducing capacity per PP was found to be 6-fold lower for the R7-knob Ad than for Ad-WTFib.

Receptor-binding capacity of recombinant virions.

Ad-FibR7-knob was assayed for its receptor binding capacity, as compared to WT virus. Both WT Ad5 and Ad-FibR7-knob virions were found to react with anti-knob mAb in ELISA, suggesting that the knob epitopes were accessible to mAb on both types of virions. However, the knob immunoreactivity was higher for WT Ad than for Ad-FibR7-knob. This suggested that the cell binding determinants of virions were more efficient or/and had a higher affinity when the knob was carried by a 22-shaft-repeat fiber, as in WT Ad5, than by a shorter, 7-repeat fiber.

These results show that the ligands which were found to be improperly folded within the cytosol and the nucleus all had a conformation that was highly dependent on the formation of disulfide bridges. Thus antibodies or fragments thereof should be used which lack disulfide bridges while retaining native or modified epitope binding.

The number of repeats in the shaft portion should preferably be at least about 7, e.g. 6-12. The presence of a wild type knob (e.g. as one of two types of fibre proteins wherein the second is modified for re-targeting) may be beneficial for generation of recombinant viable Ad virions.

- 84 -

Sequence Listing

[SEQ. ID. NO.: 1] - Adenovirus 5 Fiber Protein encoding sequence.

5

Sequence Range: 1 to 1746

```

10      10      20      30      40      50
      atgaagcgcg caagaccgtc tgaagatacc ttcaaccccg tgtatccata
10      tacttcgcgc gttctggcag acttctatgg aagttggggc acataggtat

      60      70      80      90     100
      tgacacggaa accggtcctc caactgtgcc ttttcttact cctccctttg
15      actgtgcctt tggccaggag gttgacacgg aaaagaatga ggagggaac

      110     120     130     140     150
      tatcccccaa tgggtttcaa gagagtcccc ctgggggtact ctctttgcgc
      ataggggggtt acccaaagtt ctctcagggg gaccccatga gagaaacgcg

20      160     170     180     190     200
      ctatccgaac ctctagttac ctccaatggc atgcttgccg tcaaaatggg
      gataggcttg gagatcaatg gaggttaccg tacgaacgcg agttttaccc

      210     220     230     240     250
25      caacggcctc tctctggacg aggccggcaa ccttacctcc caaaatgtaa
      gttgccggag agagacctgc tccggccggt ggaatggagg gttttacatt

      260     270     280     290     300
30      ccactgtgag cccacctctc aaaaaaacca agtcaaacad aaacctggaa
      ggtgacactc ggggtggagag tttttttggt tcagtttgta tttggacctt

      310     320     330     340     350
      atatctgcac ccctcacagt tacctcagaa gccctaactg tggctgccgc
35      tatagacgtg gggagtgtca atggagtctt cgggattgac accgacggcg

      360     370     380     390     400
      cgcacctcta atggtcgcgg gcaacacact caccatgcaa tcacaggccc
```

- 85 -

gcgtggagat taccagcgcc cgttgtgtga gtggtacgtt agtgtccggg

410 420 430 440 450
cgctaaccgt gcacgactcc aaacttagca ttgccaccca aggaccctc
5 gcgattggca cgtgctgagg tttgaatcgt aacggtgggt tcctggggag

460 470 480 490 500
acagtgtcag aaggaaagct agccctgcaa acatcaggcc ccctcaccac
10 tgtcacagtc ttcctttcga tcgggacgtt tgtagtcggg gggagtgggtg

510 520 530 540 550
caccgatagc agtaccctta ctatcactgc ctcaccccct ctaactactg
gtggctatcg tcatgggaat gatagtgcg gagtggggga gattgatgac

560 570 580 590 600
ccactggtag cttgggcatt gacttgaaag agcccattta tacacaaaat
15 ggtgaccatc gaaccogtaa ctgaactttc tcgggtaa atgtgtttta

610 620 630 640 650
ggaaaactag gactaaagta cggggctcct ttgcatgtaa cagacgacct
20 ccttttgatc ctgatttcac gccccgagga aacgtacatt gtctgctgga

660 670 680 690 700
aaacactttg accgtagcaa ctgggtccagg tgtgactatt aataatactt
25 tttgtgaaac tggcatcggt gaccagggtc acactgataa ttattatgaa

710 720 730 740 750
ccttgcaaac taaagtact ggagccttgg gttttgattc acaaggcaat
ggaacgtttg atttcaatga cctcgggaacc caaaactaag tgttccgtta

760 770 780 790 800
atgcaactta atgtagcagg aggactaagg attgattctc aaaacagacg
30 tacgttgaat tacatcggtc tcctgattcc taactaagag ttttgtctgc

810 820 830 840 850
ccttatactt gatgttagtt atccgtttga tgctcaaac caactaaatc
35 ggaatatgaa ctacaatcaa taggcaaact acgagttttg gttgatttag

- 86 -

860 870 880 890 900
taagactagg acagggccct ctttttataa actcagccca caacttggat
attctgatcc tgtcccggga gaaaaatatt tgagtcgggt gttgaaccta

5 910 920 930 940 950
attaactaca acaaaggcct ttacttgttt acagcttcaa acaattccaa
taattgatgt tgtttccgga aatgaacaaa tgcgaagtt tgttaagggt

10 960 970 980 990 1000
aaagcttgag gttaacctaa gcactgccaa ggggttgatg tttgacgcta
tttcgaactc caattggatt cgtgacgggt cccaactac aaactgcat

15 1010 1020 1030 1040 1050
cagccatagc cattaatgca ggagatgggc ttgaatttgg ttcacctaata
gtcggtatcg gtaattacgt cctctaccg aacttaaacc aagtggatta

20 1060 1070 1080 1090 1100
gcaccaaaca caaatcccct caaaacaaaa attggccatg gcctagaatt
cgtgggtttgt gtttagggga gttttgtttt taaccgggtac cggatcttaa

25 1110 1120 1130 1140 1150
tgattcaaac aaggctatgg ttcctaaact aggaactggc cttagttttg
actaagtttg ttccgatacc aaggatttga tccttgaccg gaatcaaac

30 1160 1170 1180 1190 1200
acagcacagg tgccattaca gtaggaaaca aaaataatga taagctaact
tgtcgtgtcc acggtaatgt catcctttgt ttttattact attcgattga

35 1210 1220 1230 1240 1250
ttgtggacca caccagctcc atctcctaac ttagactaa atgcagagaa
aacacctggt gtggtcgagg tagaggattg acatctgatt tacgtctctt

1260 1270 1280 1290 1300
agatgctaaa ctacttttg tcttaacaaa atgtggcagt caaatacttg
tctacgattt gagtgaacc agaattgttt tacaccgtca gtttatgaac

1310 1320 1330 1340 1350
ctacagtttc agttttggct gttaaaggca gtttggctcc aatatctgga

- 87 -

gatgtcaaag tcaaaaccga caatttccgt caaaccgagg ttatagacct

1360 1370 1380 1390 1400
acagttcaaa gtgctcatct tattataaga ttgacgaaa atggagtgt
5 tgtcaagttt cacgagtaga ataatatctt aaactgcttt tacctcacga

1410 1420 1430 1440 1450
actaaacaat tccttcctgg acccagaata ttggaacttt agaaatggag
10 tgatttgta aggaaggacc tgggtcttat aaccttgaaa tctttacctc

1460 1470 1480 1490 1500
atcttactga aggcacagcc tatacaaacg gtgttggatt tatgcctaac
tagaatgact tccgtgtcgg atatgtttgc cacaacctaa atacggattg

1510 1520 1530 1540 1550
ctatcagctt atccaaaatc tcacggtaaa actgccaaaa gtaacattgt
gatagtcgaa taggttttag agtgccattt tgacggtttt cattgtaaca

1560 1570 1580 1590 1600
cagtcaagtt tacttaaacg gagacaaaac taaacctgta acactaacca
20 gtcagttcaa atgaatttgc ctctgttttg atttggacat tgtgattggt

1610 1620 1630 1640 1650
ttacactaaa cggtaacacg gaaacaggag acacaactcc aagtgcatac
25 aatgtgattt gccatgtgtc ctttgtcctc tgtgttgagg ttcacgtatg

1660 1670 1680 1690 1700
tctatgtcat tttcatggga ctggtctggc cacaactaca ttaatgaaat
agatacagta aaagtaccct gaccagaccg gtgttgatgt aattacttta

30
1710 1720 1730 1740
atttgccaca tcctcttaca ctttttcata cattgcccaa gaataa
taaacggtgt aggagaatgt gaaaaagtat gtaacgggtt cttatt

- 88 -

[SEQ. ID. NO.: 2] - Neck Region Peptide from Human Lung Surfactant Protein D.

PDVASLRQQVEALQGQVQHLQAAFSQYKKVELFPNG

5

[SEQ. ID. NO.: 3] - Recombinant Fiber A1

gaattcg atg aag cgc gca aga ccg tct gaa gat acc ttc aac ccc gtg tat
cca tat gac acg gaa acc ggt cct cca act gtg cct ttt ctt act cct ccc
10 ttt gta tcc ccc aat ggg ttt caa gag agt ccc cct ggg gta ctc tct ttg
cgc cta tcc gaa cct cta gtt acc tcc aat ggc atg cct gac gta gca agc
tta cga caa cag gta gaa gcc ttg caa ggg cag gta caa cac tta cag gcg
gca ttt agc caa tac aaa aag gta gag ttg ttt cca aac gga gcc aag aag
ctg aac gac gcc cag gcc ccc aag agc gac cca tcg atc taa ctc gag
15

[SEQ. ID. NO.: 4] - Human epidermal growth factor (EGF)

Sequence Range: 1 to 183

20

10 20 30 40 50
gggaattcat gaactccgac tccgaatgtc cattgtccca cgacgggttac
cccttaagta cttgaggctg aggcttacag gtaacagggt gctgccaatg
M N S D S E C P L S H D G Y>

25

60 70 80 90 100
tgtttgcacg acggtgtttg tatgtacatc gaagctttgg acaagtacgc
acaaacgtgc tgccacaaac atacatgtag cttcgaaacc tgttcatgcg
C L H D G V C M Y I E A L D K Y A>

30

110 120 130 140 150
ttgtaactgt gttgttggtt acatcggtga aagatgtcaa tacagagact
aacattgaca caacaaccaa thtagccact ttctacagtt atgtctctga
35 C N C V V G Y I G E R C Q Y R D>

- 89 -

160 170 180
tgaagtgggtg ggaattgaga tgataagaat tcc
acttcaccac ccttaactct actattctta agg
L K W W E L R *>

5

[SEQ. ID. NO.: 5] - Amino acid linker from Staphylococcal protein A

10 Sequence Range: 1 to 39

10 20 30
gccaagaagc tgaacgacgc ccaggccccc aagagcgac
cggttcttcg acttgctgcg ggtccggggg ttctcgctg
15 A K K L N D A Q A P K S D>
_____a_____SPA 48-60_____a_____

[SEQ. ID. NO.: 6] - Fiber construct A1 EGF

20 Sequence Range: 1 to 513

10 20 30 40 50
atgaagcgcg caagaccgtc tgaagatacc ttcaaccccg tgtatccata
tacttcgcgc gttctggcag acttctatgg aagttggggc acataggtat
25 M K R A R P S E D T F N P V Y P Y>
_____b_____b_TAIL_____b_____b_____>

60 70 80 90 100
tgacacggaa accggtcctc caactgtgcc ttttcttact cctcccttg
actgtgcctt tggccaggag gttgacacgg aaaagaatga ggagggaaac
30 D T E T G P P T V P F L T P P F>
_____b_____b_TAIL_____b_____b_____>

110 120 130 140 150
tatcccccaa tgggtttcaa gagagtcccc ctgggggtact ctctttgcgc
35 ataggggggtt acccaaagtt ctctcagggg gaccccatga gagaaacgcg
L S L R>

- 90 -

_____a_____>
V S P N G F Q E S P P G V>
_____b_____TAIL_____b_____>

5 160 170 180 190 200
ctatccgaac ctctagttac ctccaatggc atgcctgacg tagcaagctt
gataggcttg gagatcaatg gaggttaccg tacggactgc atcgttcgaa
L S E P L V T S N G M>
_____a_____R1_____a_____a_____>

10 P D V A S L>
 ____LUNG SURFAC____>

 210 220 230 240 250
acgacaacag gtagaagcct tgcaagggca ggtacaacac ttacaggcgg
15 tgctgttgtc catcttcgga acgttcccgt ccatgttggtg aatgtccgcc
R Q Q V E A L Q G Q V Q H L Q A>
_____LUNG SURFACTANT PROTEIN D TRIM_____c_____>

 260 270 280 290 300
catttagcca atacaaaaag gtagagttgt ttccaaacgg agccaagaag
20 gtaaatecgt tatgtttttc catctcaaca aaggtttgcc tcggttcttc
A F S Q Y K K V E L F P N G>
_____LUNG SURFACTANT PROTEIN D TRIM_____c_____>
 A K K>
25 _____>

 >Cla1
 |
 310 320 330 340 350
30 ctgaacgacg cccaggcccc caagagcgac ccatcgatca tgaactccga
gacttgctgc ggggtccgggg gttctcgctg ggtagctagt acttgaggct
M N S D>
_____d_____>

 L N D A Q A P K S D>
35 SPA 48-60_e_____>
 P S I>
 _____>

- - 91 - -

```

                    360      370      380      390      400
ctccgaatgt ccattgtccc acgacgggta ctgtttgcac gacgggtgtt
gaggcttaca ggtaacaggg tgctgccaat gacaaacgtg ctgccacaaa
5      S E C P L S H D G Y C L H D G V>
___EPIDERMAL GROWTH FACTOR (UROGASTRONE); CODON_ST___>

                    410      420      430      440      450
gtatgtacat cgaagctttg gacaagtacg cttgtaactg tgttgttggt
10     catacatgta gcttcgaaac ctgttcattgc gaacattgac acaacaacca
      C M Y I E A L D K Y A C N C V V G>
___EPIDERMAL GROWTH FACTOR (UROGASTRONE); CODON_ST___>

                    460      470      480      490      500
15     tacatcggtg aaagatgtca atacagagac ttgaagtggg gggaattgag
      atgtagccac tttctacagt tatgtctctg aacttcacca cccttaactc
      Y I G E R C Q Y R D L K W W E L R>
___EPIDERMAL GROWTH FACTOR (UROGASTRONE); CODON_ST___>

20     >xho1
      |
      | 510
      atgactcgag ggg
      tactgagctc ccc
25     X>
      ___>

```

[SEQ. ID. NO.: 7] - Fiber Construct A7 EGF

30 Sequence Range: 1 to 811

```

>EcoR1
|
| 10      20      30      40      50
35     gaattcgatg aagcgcgcaa gaccgtctga agataccttc aaccccgtgt

```

- - 92 - -

cttaagctac ttcgcgcggtt ctggcagact tctatggaag ttggggcaca
 M>
 ____>
 M K R A R P S E D T F N P V>
 5 ____j____j____TAIL____j____>

60 70 80 90 100
 atccatatga cacggaaacc ggtcctccaa ctgtgccttt tcttactcct
 taggtatact gtgcctttgg ccaggagggtt gacacggaaa agaagtagga
 10 Y P Y D T E T G P P T V P F L T P>
 ____j____j____TAIL____j____j____>

110 120 130 140 150
 ccctttgtat cccccaatgg gtttcaagag agtccccctg gggtagctctc
 15 gggaaacata ggggggttacc caaagttctc tcagggggac cccatgagag
 L S>
 ____>

P F V S P N G F Q E S P P G V>
 ____j____TAIL____j____j____>

20 160 170 180 190 200
 tttgcgccta tccgaacctc tagttacctc caatggcatg cttgcgctca
 aaacgcggat aggcttgag atcaatggag gttaccgtac gaacgcgagt
 L A L>
 ____>

L R L S E P L V T S N G M>
 ____h____REPEAT 1____h____>

210 220 230 240 250
 30 aaatgggcaa cggcctctct ctggacgagg ccggcaacct tacctcccaa
 tttaccggtt gccggagaga gacctgctcc ggccgttgga atggagggtt
 L T S Q>
 ____f____>

K M G N G L S L D E A G N>
 35 ____g____REPEAT 2____g____>

- - 93 - -

260 270 280 290 300
 aatgtaacca ctgtgagccc acctctcaaa aaaaccaagt caaacataaa
 ttacattggg gacactcggg tggagagttt ttttggttca gtttgatatt
 I N>
 5 _____>
 N V T T V S P P L K K T K S N>
 _____f_____REPEAT 3 _____f_____f_____>
 310 320 330 340 350
 10 cctggaaata tctgcacccc tcacagttac ctcaagaagcc ctaactgtgg
 ggacctttat agacgtgggg agtgtcaatg gagtcttcgg gattgacacc
 L T V>
 _____>
 L E I S A P L T V T S E A>
 15 _____e_____REPEAT 4 _____e_____>
 360 370 380 390 400
 ctgccgccgc acctctaattg gtcgcgggca acacactcac catgcaatca
 gacggcggcg tggagattac cagcgcccgt tgtgtgagtg gtacgttagt
 20 L T M Q S>
 _____REPEAT 6_____>
 A A A A P L M V A G N T>
 _____d_____REPEAT 5 _____d_____>
 410 420 430 440 450
 25 caggccccgc taaccgtgca cgactccaaa cttagcattg ccaccaagg
 gtccggggcg attggcacgt gctgaggttt gaatcgtaac ggtgggttcc
 L S I A T Q G>
 _____REPEAT 7_____>
 30 Q A P L T V H D S K>
 _____REPEAT 6_c_____>
 460 470 480 490 500
 35 acccctcaca gtgtcagaag gaaagctagc ccctgacgta gcaagcttac
 tggggagtggt cacagtcttc ctttcgatcg gggactgcat cgttcgaatg

- - 94 - -

5 L A>
_____a_>
P L T V S E G K>
_____REPEAT 7_____b_____>

510 520 530 540 550
gacaacaggt agaagccttg caagggcagg tacaacactt acagggcggca
ctgttgtcca tcttcggaac gttcccgtcc atgttgtgaa tgtccgccgt
10 R Q Q V E A L Q G Q V Q H L Q A A>
_____LUNG SURFACTANT PROTEIN D TRIM_____k_____>

560 570 580 590 600
tttagccaat acaaaaaggt agagttgttt ccaaacggag ccaagaagct
15 aaatcgggta tgtttttcca tctcaacaaa ggtttgcctc ggttcttcga
F S Q Y K K V E L F P N G>
_____LUNG SURFACTANT PROTEIN D TRIM_____>

_____A K K L>
_____m_____>

20 >Cla1
|

610 620 630 | 640 650
gaacgacgcc caggccccca agagcgaccc atcgatcatg aactccgact
25 cttgctgcgg gtccgggggt tctcgctggg tagctagtac ttgaggctga
M N S D>
_____l_____>

N D A Q A P K S D>
_____SPA 48-60_____m_____>

30 P S I>
_____n_____>

660 670 680 690 700
ccgaatgtcc attgtccac gacggttact gtttgcacga cgggtgtttgt
35 ggcttacagg taacaggggtg ctgccaatga caaacgtgct gccacaaaca

- - 95 - -

S E C P L S H D G Y C L H D G V C>
____EPIDERMAL GROWTH FACTOR (UROGASTRONE); CODON_ST____>

710 720 730 740 750
5 atgtacatcg aagctttgga caagtacgct tgtaactgtg ttgttggtta
tacatgtagc ttcgaaacct gttcatgcga acattgacac aacaaccaat
M Y I E A L D K Y A C N C V V G Y>
____EPIDERMAL GROWTH FACTOR (UROGASTRONE); CODON_ST____>
760 770 780 790 800
10 catcgggtgaa agatgtcaat acagagactt gaagtgggtgg gaattgagat
gtagccactt tctacagtta tgtctctgaa cttcaccacc cttaactcta
I G E R C Q Y R D L K W W E L R>
____EPIDERMAL GROWTH FACTOR (UROGASTRONE); CODON_ST____>

15 >xho1
|
| 810
gactcgaggg g
ctgagctccc c
20 X>
_>

[SEQ. ID. NO.: 8] - G250 Construct

25 Sequence Range: 1 to 1056

10 20 30 40 50
gacattgtga tgaccagtc tcaaagattc atgtccacaa cagtaggaga
ctgtaacact actgggtcag agtttctaag tacagggtgtt gtcacacctt
30 D I V M T Q S Q R F M S T T V G D>
____c____c____VK____c____c____>
60 70 80 90 100
cagggtcagc atcacctgca aggccagtca gaatgtggtt tctgctgttg
35 gtcccagtcg tagtgagcgt tccgggtcagt cttacaccaa agacgacaac

- - 96 - -

R V S I T C K A S Q N V V S A V>
_____c_____c_VK_____c_____c_____>

110 120 130 140 150
5 cctggatatca acagaaacca ggacaatctc cttaaactact gatttactca
ggaccatagt tgtcttttggc cctggttagag gatttgatga ctaaatgagt
A W Y Q Q K P G Q S P K L L I Y S>
_____c_____c_VK_____c_____c_____>

160 170 180 190 200
10 gcatccaatc ggtacactgg agtccctgat cgcttcacag gcagtggatc
cgtaggtag ccatgtgacc tcagggacta gcgaagtgtc cgtcacctag
A S N R Y T G V P D R F T G S G S>
_____c_____c_VK_____c_____c_____>

210 220 230 240 250
15 tgggacagat ttcactctca ccattagcaa tatgcagtct gaagacctgg
accctgtcta aagtgagagt ggtaatcggt atacgtcaga cttctggacc
G T D F T L T I S N M Q S E D L>
_____c_____c_VK_____c_____c_____>

260 270 280 290 300
20 ctgatttttt ctgtcaacaa tatagcaact atccgtggac gttecggtgga
gactaaaaaa gacagttggt atatcggtga taggcacctg caagccacct
A D F F C Q Q Y S N Y P W T F G G>
_____c_____c_VK_____c_____c_____>

310 320 330 340 350
25 ggcaccaagc tggaaatcaa aggatctggc tctacttccg gtagcggcaa
ccgtgggttcg acctttagtt tcctagaccg agatgaaggc catcgccgtt
G T K L E I K>
_____VK_____c_____>

310 320 330 340 350
30 ggcaccaagc tggaaatcaa aggatctggc tctacttccg gtagcggcaa
ccgtgggttcg acctttagtt tcctagaccg agatgaaggc catcgccgtt
G T K L E I K>
_____VK_____c_____>

310 320 330 340 350
35 G S G S T S G S G K>
_____d_LINK_____d_____>

- - 97 - -

360 370 380 390 400
atcctctgaa ggcaaaggta ctagagacgt gaagctcgtg gagtctgggg
taggagactt ccgtttccat gatctctgca cttcgagcac ctcagacccc
D V K L V E S G>
5 _____a_____VH_____a_____>
S S E G K G T R>
_____LINK_____d_____>

410 420 430 440 450
10 gaggcttagt gaagcttgga gggccctga aactctcctg tgcagcctct
ctccgaatca cttcgaacct cccagggact ttgagaggac acgtcggaga
G G L V K L G G S L K L S C A A S>
_____a_____a_____VH_____a_____a_____>

460 470 480 490 500
15 ggattcactt tcagtaacta ttacatgtct tgggttcgcc agactccaga
cctaagtga agtcattgat aatgtacaga acccaagcgg tctgaggtct
G F T F S N Y Y M S W V R Q T P E>
_____a_____a_____VH_____a_____a_____>

510 520 530 540 550
20 gaagaggctg gagttggctg cagccattaa tagtgatggt ggtatcacct
cttctccgac ctcaaccagc gtcggtaatt atcactacca ccatagtgga
K R L E L V A A I N S D G G I T>
25 _____a_____a_____VH_____a_____a_____>

560 570 580 590 600
actatctaga cactgtgaag ggccgattca ccatttcaag agacaatgcc
tgatagatct gtgacacttc ccggctaagt ggtaaagttc tctgttacgg
30 Y Y L D T V K G R F T I S R D N A>
_____a_____a_____VH_____a_____a_____>

610 620 630 640 650
35 aagaacaccc tgtacctgca aatgagcagt ctgaagtctg aggacacagc
ttcttgggg acatggacgt ttactcgtca gacttcagac tcctgtgtcg

```

        660             670             680             690             700
cttgtttttac tgtgcaagac accgctcggg ctacttttct atggactact
gaacaaaatg acacgttctg tggcgagccc gatgaaaaga tacctgatga
  L  F  Y   C  A  R   H  R  S  G   Y  F  S   M  D  Y>
          a             a  VH             a             a             >

```

15

A>

_____>

C P P C P>

HINGE >

25

810 820 830 840 850

ggacaccctc atgatctccc ggaccacctga ggtcacatgc gtgggtgggtgg
cctgtggggag tactagaggg cctgggggact ccagtgtacg caccaccacc

D T L M I S R T P E V T C V V V>

30

b b CH2 b b

```

                860                870                880                890                900
acgtgagcca cgaagaccct gaggtcaagt tcaactggta cgtggacggc
tgcaactcggg gcttctggga ctccagttca agttgaccat gcacctgccg
35 D V S H E D P E V K F N W Y V D G>

```

- - 99 - -

```

_____b_____b___CH2___b_____b_____>

          910          920          930          940          950
gtggaggtgc ataatgccaa gacaaagccg cgggaggagc agtacaacag
5  cacctccacg tattacgggt ctgtttcggc gccctcctcg tcatgttgtc
   V E V H N A K T K P R E E Q Y N S>
_____b_____b___CH2___b_____b_____>

          960          970          980          990          1000
10  cacgtaccgg gtggtcagcg tcctcaccgt cctgcaccag gactggctga
   gtgcatggcc caccagtcgc aggagtggca ggacgtggtc ctgaccgact
   T Y R V V S V L T V L H Q D W L>
_____b_____b___CH2___b_____b_____>

          1010          1020          1030          1040          1050
15  atggcaagga gtacaagtgc aaggtctcca acaaagccct cccagccccc
   taccgttcct catgttcacg ttccagaggt tgtttcggga gggtcggggg
   N G K E Y K C K V S N K A L P A P>
_____b_____b___CH2___b_____b_____>

20  atcgag
   tagctc
   I E>
_____>

25

```

[SEQ. ID. NO.: 9] - C242 Construct

30 Sequence Range: 1 to 793

```

          >Sfi          >EcoR1          >ClaI
          |          |          |
          |          |          |
          |    10    |    20    |    30          40          50
35  gcggcccagc cggccacgaa ttcatcgatg gatattgtga tgactcaggc

```

- - 100 - -

cgccggggtcg gccgggtgctt aagtagctac ctataacact actgagtcgg

A Q P A T N S S M>

_____b BACKPRIMER_____>

D I V M T Q A>

5

_____VK PHARMACIA_____>

A>

____>

10

60 70 80 90 100
tgcaccctct gtacctgtca ctctggaga gtcagtatcc atctcctgca
acgtgggaga catggacagt gaggacctct cagtcatagg tagaggacgt

A P S V P V T P G E S V S I S C>

_____d_____VK PHARMACIA_____d_____>

15

110 120 130 140 150
ggtctagtaa gagtctcctg catagtaatg gcaacactta cttgtattgg
ccagatcatt ctcagaggac gtatcattac cgttgtgaat gaacataacc
R S S K S L L H S N G N T Y L Y W>

_____d_____VK PHARMACIA_____d_____>

20

S L L H S N G N T Y L Y>

_____f_____CDR1_____f_____>

25

160 170 180 190 200
ttcctgcaga ggccaggcca gtctcctcag ctctgatat atcggatgtc
aaggacgtct ccggtccggt cagaggagtc gaggactata tagcctacag
F L Q R P G Q S P Q L L I Y R M S>

_____d_____VK PHARMACIA_____d_____>

R M S>

30

210 220 230 240 250
caaccttgctc tcaggagtcc cagacagggt cagtggcagt gggtcaggaa
gttggaacag agtcctcagg gtctgtccaa gtcaccgtca cccagtcctt
N L V S G V P D R F S G S G S G>

35

_____d_____VK PHARMACIA_____d_____>

- - 101 - -

N L V S>

_____e____>

260 270 280 290 300
5 ctgctttcac actgagaatc agtagagtgg aggctgagga tgtgggtgtt
gacgaaagtg tgactcttag tcatctcacc tccgactcct acacccacaa
T A F T L R I S R V E A E D V G V>
_____d_____VK PHARMACIA_____d_____>

310 320 330 340 350
10 tattactgtc tgcaacatct agagtatccg ttcacgttcg gtcctgggac
ataatgacag acgttgtaga tctcataggc aagtgcgaagc caggaccctg
L Q H L E Y P F T>
_a_____CDR3_____a_____>
15 Y Y C L Q H L E Y P F T F G P G T>
_____d_____VK PHARMACIA_____d_____>

360 370 380 390 400
20 caagctggag ctgaaacggc ccccgactt tggtcccccg gccgctagtt
gttcgacctc gactttgccg ggggcctgaa acaagggggc cggcgatcaa
K L E L K R>
___VK PHARMACIA___>
P P D F V P P A A S>
_1_____1_825_____1_____>

410 420 430 440 450
25 tccctgatca ctcccctcgt ggccagggtcc agttggtgca gtctggacct
agggactagt gaggggagca ccggtccagg tcaaccacgt cagacctgga
Q V Q L V Q S G P>
30 _____VH PHARMACIA [SPLIT]_____>
F P D H S P R G>
_____825_____1_____>

460 470 480 490 500
35 gagctgaaga agcctggaga gacagtcaag atctcctgca aggcttctga

- - 102 - -

ctcgacttct tcggacctct ctgtcagttc tagaggacgt tccgaagact
E L K K P G E T V K I S C K A S D>
_____h_____VH PHARMACIA [SPLIT]_____h_____>

5 510 520 530 540 550
ttataccttc acatactatg gaatgaactg ggtgaagcag gctccgggaa
aatatggaag tgtatgatac cttacttgac ccacttcgtc cgaggccctt
Y T F T Y Y G M N W V K Q A P G>
_____h_____VH PHARMACIA [SPLIT]_____h_____>

10 Y Y G M N>
 _____CDR1_____>

 560 570 580 590 600
agggttttaa gtggatgggc tggatagaca ccaccactgg agagccaaca
tcccaaattt cacctacccg acctatctgt ggtggtgacc tctcggttgt
K G L K W M G W I D T T T G E P T>
_____h_____VH PHARMACIA [SPLIT]_____h_____>
 W I D T T T G E P T>
 _____CDR2 [SPLIT]_____>

20

 610 620 630 640 650
tatgctgaag attttaaggg acggattgcc ttctctttgg agacctctgc
atacgacttc taaaattccc tgcctaacgg aagagaaacc tctggagacg
Y A E D F K G R I A F S L E T S A>
_____h_____VH PHARMACIA [SPLIT]_____h_____>
Y A E D F K G>
_____CDR2 [SPLIT]_____i_>

25

 660 670 680 690 700
cagcactgcc tatttgcaga tcaaaaacct caaaaatgag gacacggcta
gtcgtgacgg ataaacgtct agttttttgga gtttttactc ctgtgccgat
S T A Y L Q I K N L K N E D T A>
_____h_____VH PHARMACIA [SPLIT]_____h_____>

30

35 710 720 730 740 750

catatttctg tgcaagacgg gggccttaca actggtactt tgatgtctgg
gtataaagac acgttctgcc cccggaatgt tgaccatgaa actacagacc

_____g_____CDR3_____g_____>

T Y F C A R R G P Y N W Y F D V W>

```
>Xho(-)StopXho_adapt_[Split]
```

790

ccggttcctt ggtgccagt ggcagaggagt gagctaattg agc

_____ >

G Q G T T V T V S S>

 VH PHARMACIA [SPLIT] >

5

1

10

 >

15

60 70 80 90 100

20

110 120 130 140 150

25

_____ >

_____f_____TAIL_____f_____f_____>

160 170 180 190 200

30

L R L S E P L V T S N G M>

35

BNSDOCID: <WO_____0208263A2 | >

- - 105 - -

_____>

210 220 230 240 250
caagcttacg acaacaggta gaagccttgc aagggcaggt acaacactta
5 gttcgaatgc tgttgtccat ctccggaacg ttcccgtcca tgttgtgaat
A S L R Q Q V E A L Q G Q V Q H L>
_____LUNG SURFACTANT PROTEIN D TRIM_g_____>

260 270 280 290 300
10 caggcggcat ttagccaata caaaaaggta gagttgtttc caaacggagc
gtccgccgta aatcggttat gtttttccat ctcaacaaag gtttgcctcg
A>
____>

Q A A F S Q Y K K V E L F P N G>
15 _____LUNG SURFACTANT PROTEIN D TRIM_g_____>

>Cla1
|

310 320 330 340 350
20 caagaagctg aacgacgccc agggcccca gagcgacca tcgatcgaca
gttcttcgac ttgctgcggg tccgggggtt ctgctgggt agctagctgt
K K L N D A Q A P K S D>
_____b_____SPA 48-60_____b_____>

P S I>
25 _____c_____>
D>
_____>

360 370 380 390 400
30 ttgtgatgac ccagtctcaa agattcatgt ccacaacagt aggagacagg
aacactactg ggtcagagtt tctaagtaca ggtgttgtca tcctctgtcc
I V M T Q S Q R F M S T T V G D R>
_____j_____j_____VK_____j_____>

410 420 430 440 450
35

- - 106 - -

gtcagcatca cctgcaaggc cagtcagaat gtggtttctg ctgttgccctg
cagtcgtagt ggacgttccg gtcagtctta caccaaagac gacaacggac
V S I T C K A S Q N V V S A V A W>
_____j_____j_____VK_____j_____j_____>
460 470 480 490 500

5
gtatcaacag aaaccaggac aatctcctaa actactgatt tactcagcat
catagttgtc tttggtcctg ttagaggatt tgatgactaa atgagtcgta
Y Q Q K P G Q S P K L L I Y S A>
_____j_____j_____VK_____j_____j_____>

10
510 520 530 540 550
ccaatcggta cactggagtc cctgatcgct tcacaggcag tggatctggg
ggtagccat gtgacctcag ggactagcga agtgtccgct acctagaccc
S N R Y T G V P D R F T G S G S G>
_____j_____j_____VK_____j_____j_____>

15
560 570 580 590 600
acagatttca ctctcaccat tagcaatatg cagtctgaag acctggctga
tgtctaaagt gagagtggta atcggtatac gtcagacttc tggaccgact
T D F T L T I S N M Q S E D L A D>
_____j_____j_____VK_____j_____j_____>

20
610 620 630 640 650
ttttttctgt caacaatata gcaactatcc gtggacgttc ggtggaggca
aaaaaagaca gttgttatat cggtgatagg cacctgcaag ccacctccgt
F F C Q Q Y S N Y P W T F G G G>
_____j_____j_____VK_____j_____j_____>

25
660 670 680 690 700
ccaagctgga aatcaaagga tctggctcta cttccggtag cggcaaattc
ggttcgacct ttagtttctt agaccgagat gaaggccatc gccgtttagg
T K L E I K>
_____VK_j_____>

30
G S G S T S G S G K S>
_____k_____LINK_____k_____>

35

- - 107 - -

710 720 730 740 750
tctgaaggca aaggtactag agacgtgaag ctctgtggagt ctggggggagg
agacttccgt ttccatgata tctgcacttc gagcacctca gacccccctcc
D V K L V E S G G G>
5 _____h_____VH_____h_____>
S E G K G T R>
_____LINK_____k_>

760 770 780 790 800
10 cttagtgaag cttggagggt ccctgaaact ctctgtgca gcctctggat
gaatcacttc gaacctcca gggactttga gaggacacgt cggagaccta
L V K L G G S L K L S C A A S G>
_____h_____h_____VH_____h_____h_____>

810 820 830 840 850
15 tcactttcag taactattac atgtcttggg ttccgacagac tccagagaag
agtgaagtc attgataatg tacagaaccc aagcggctctg aggtctcttc
F T F S N Y Y M S W V R Q T P E K>
_____h_____h_____VH_____h_____h_____>

860 870 880 890 900
20 aggctggagt tggctgcagc cattaatagt gatgggtggta tcacctacta
tccgacctca accagcgtcg gtaattatca ctaccaccat agtggatgat
R L E L V A A I N S D G G I T Y Y>
25 _____h_____h_____VH_____h_____h_____>

910 920 930 940 950
30 tctagacact gtgaagggcc gattcaccat ttcaagagac aatgccaaga
agatctgtga cacttcccgg ctaagtggta aagttctctg ttacggttct
L D T V K G R F T I S R D N A K>
_____h_____h_____VH_____h_____h_____>

960 970 980 990 1000
35 acaccctgta cctgcaaata agcagtctga agtctgagga cacagccttg
tgtgggacat ggacgtttac tcgtcagact tcagactcct gtgtcggaac

- - 108 - -

5

10

15

20

25

30

35

- - 109 - -

```
_____i_____i_____CH2_____i_____i_____>

      1260      1270      1280      1290      1300
aggtgcataa tgccaagaca aagccgcggg aggagcagta caacagcacg
5 tccacgtatt acggttctgt ttcggcgccc tctcgtcat gttgtcgtgc
  E V H N A K T K P R E E Q Y N S T>
_____i_____i_____CH2_____i_____i_____>

      1310      1320      1330      1340      1350
10 taccgggtgg tcagcgtcct caccgtcctg caccaggact ggctgaatgg
  atggcccacc agtcgcagga gtggcaggac gtggtcctga ccgacttacc
  Y R V V S V L T V L H Q D W L N G>
_____i_____i_____CH2_____i_____i_____>

      1360      1370      1380      1390      1400
15 caaggagtac aagtgcaagg tctccaacaa agccctccca gcccccatcg
  gttcctcatg ttcacgttcc agaggttggt tcgggagggt cgggggtagc
  K E Y K C K V S N K A L P A P I>
_____i_____i_____CH2_____i_____i_____>

20
    >XhoI
      |
      |1410
agtaactcga g
25 tcattgagct c
    * L E>
      _____a_>
    E>
    ____>

30
```

[SEQ. ID. NO.: 11] - Fiber Construct A7 G250

35 Sequence Range: 1 to 1702

- - 110 - -

>EcoR1

5
|
| 10 20 30 40 50
gaattcgatg aagcgcgcaa gaccgtctga agataccttc aaccccggtgt
cttaagctac ttcgcgcggtt ctggcagact tctatggaag ttggggcaca
M>
____>
M K R A R P S E D T F N P V>
____m____m____TAIL____m____>
10
60 70 80 90 100
atccatatga cacggaacc ggtcctccaa ctgtgccttt tcttactcct
taggtatact gtgccttttg ccaggagggtt gacacggaaa agaatzgagga
Y P Y D T E T G P P T V P F L T P>
15
____m____m____TAIL____m____m____>
110 120 130 140 150
ccctttgtat cccccaatgg gtttcaagag agtccccctg gggactctc
gggaaacata ggggggttacc caaagttctc tcagggggac cccatgagag
20
L S>
____>
P F V S P N G F Q E S P P G V>
____m____TAIL____m____m____>
25
160 170 180 190 200
tttgcgccta tccgaacctc tagttacctc caatggcatg cttgcgctca
aaacgcggat aggcttggag atcaatggag gttaccgtac gaacgcgagt
L A L>
____>
30
L R L S E P L V T S N G M>
____k____REPEAT 1____k____>
210 220 230 240 250
aaatgggcaa cggcctctct ctggacgagg ccggcaacct tacctcccaa
35
tttaccggtt gccggagaga gacctgctcc ggccgttgga atggagggtt

- - 111 - -

5
L T S Q>
____i____>
K M G N G L S L D E A G N>
____j____REPEAT 2____j____>

260 270 280 290 300
aatgtaacca ctgtgagccc acctctcaaa aaaaccaagt caaacataaa
ttacattggt gacactcggg tggagagttt ttttggttca gtttgtattt
I N>
____>

10
N V T T V S P P L K K T K S N>
____i____REPEAT 3____i____i____>
310 320 330 340 350
cctggaaata tctgcacccc tcacagttac ctcagaagcc ctaactgtgg
ggacctttat agacgtgggg agtgtcaatg gagtcttcgg gattgacacc
L T V>
____>

15
L E I S A P L T V T S E A>
____h____REPEAT 4____h____>

20
360 370 380 390 400
ctgccgccgc acctctaattg gtcgcgggca acacactcac catgcaatca
gacggcgggcg tggagattac cagcgcccgt tgtgtgagtg gtacgttagt
L T M Q S>
____REPEAT 6____>

25
A A A A P L M V A G N T>
____g____REPEAT 5____g____>

30
410 420 430 440 450
caggccccgc taaccgtgca cgactccaaa cttagcattg ccacccaagg
gtccggggcg attggcacgt gctgaggttt gaatcgtaac ggtgggttcc
L S I A T Q G>
____REPEAT 7____>

35
Q A P L T V H D S K>
____REPEAT 6____f____>

- - 112 - -

460 470 480 490 500
acccttcaca gtgtcagaag gaaagctagc ccctgacgta gcaagcttac
tggggagtggt cacagtcttc ctttcgatcg gggactgcat cgttcgaatg
L A>
5 _____d____>
P L T V S E G K>
____REPEAT 7____e____>
P D V A S L>
10 _____LUNG SURFACTA____>
510 520 530 540 550
gacaacaggt agaagccttg caagggcagg tacaacactt acaggcggca
ctgttggtcca tcttcggaac gttcccgtcc atgttggtgaa tgtccgccgt
R Q Q V E A L Q G Q V Q H L Q A A>
15 _____LUNG SURFACTANT PROTEIN D TRIM____n____>
560 570 580 590 600
tttagccaat acaaaaaggt agagttgttt ccaaacggag ccaagaagct
aaatcggtta tgtttttcca tctcaacaaa ggtttgcctc ggttcttcga
20 A K K L>
_____b____>
F S Q Y K K V E L F P N G>
____LUNG SURFACTANT PROTEIN D TRIM____>
25 >Cla1
|
610 620 630 | 640 650
gaacgacgcc caggcccca agagcgaccc atcgatcgac attgtgatga
cttgctgcgg gtccgggggt tctcgctggg tagctagctg taacactact
30 N D A Q A P K S D>
____SPA 48-60____b____>
P S I>
_____c____>
D I V M>
35 _____a____>

- - 113 - -

5 660 670 680 690 700
cccagtctca aagattcatg tccacaacag taggagacag ggtcagcatc
gggtcagagt ttctaagtac aggtgttgtc atcctctgtc ccagtcgtag
T Q S Q R F M S T T V G D R V S I>
_____q_____q_VK_____q_____q_____>

10 710 720 730 740 750
acctgcaagg ccagtcagaa tgtggtttct gctgttgccct ggtatcaaca
tggaegttcc ggtcagtctt acaccaaga cgacaacgga ccatagttgt
T C K A S Q N V V S A V A W Y Q Q>
_____q_____q_VK_____q_____q_____>

15 760 770 780 790 800
gaaaccagga caatctccta aactactgat ttactcagca tccaatcggg
ctttgggtcct gttagaggat ttgatgacta aatgagtcgt aggttagcca
K P G Q S P K L L I Y S A S N R>
_____q_____q_VK_____q_____q_____>

20 810 820 830 840 850
aactggaggt ccctgatcgc ttcacaggca gtggatctgg gacagatttc
tgtgacctca gggactagcg aagtgtccgt cacctagacc ctgtctaaag
Y T G V P D R F T G S G S G T D F>
_____q_____q_VK_____q_____q_____>

25 860 870 880 890 900
actctcacca ttagcaatat gcagtctgaa gacctggctg attttttctg
tgagagtggg aatcggtata cgtcagactt ctggaccgac taaaaaagac
T L T I S N M Q S E D L A D F F C>
_____q_____q_VK_____q_____q_____>

30 910 920 930 940 950
tcaacaatat agcaactatc cgtggacgtt cggtggaggc accaagctgg
agttgttata tcgttgatag gcacctgcaa gccacctccg tggttcgacc
Q Q Y S N Y P W T F G G G T K L>
_____q_____q_VK_____q_____q_____>

35

- - 114 - -

960 970 980 990 1000
 aaatcaaagg atctggctct acttccggta gcggcaaata ctctgaaggc
 tttagtttcc tagaccgaga tgaaggccat cgccgttttag gagacttccg
 E I K>
 5 _____>
 G S G S T S G S G K S S E G>
 _r_____r_____LINK_____r_____>
 1010 1020 1030 1040 1050
 10 aaaggtacta gagacgtgaa gctcgtggag tctggggggag gcttagtgaa
 tttccatgat ctctgcactt cgagcacctc agaccccctc cgaatcactt
 D V K L V E S G G G L V K>
 _____o_____VH_____o_____>
 K G T R>
 15 _____r_____>
 1060 1070 1080 1090 1100
 gcttggaggg tccctgaaac tctcctgtgc agcctctgga ttcactttca
 cgaacctccc agggactttg agaggacacg tcggagacct aagtgaaggt
 L G G S L K L S C A A S G F T F>
 20 _____o_____o_____VH_____o_____o_____>
 1110 1120 1130 1140 1150
 gtaactatta catgtcttgg gttcgccaga ctccagagaa gaggctggag
 cattgataat gtacagaacc caagcgggtct gaggtctctt ctccgacctc
 25 S N Y Y M S W V R Q T P E K R L E>
 _____o_____o_____VH_____o_____o_____>
 1160 1170 1180 1190 1200
 ttggtcgcag ccattaatag tgatggtggt atcacctact atctagacac
 30 aaccagcgtc ggtaattatc actaccacca tagtggtatga tagatctgtg
 L V A A I N S D G G I T Y Y L D T>
 _____o_____o_____VH_____o_____o_____>
 1210 1220 1230 1240 1250
 35 tgtgaagggc cgattcacca tttcaagaga caatgccaaag aacaccctgt

- - 115 - -

acacttcccg gctaagtggg aaagttctct gttacgggttc ttgtgggaca
V K G R F T I S R D N A K N T L>

_____o_____o_VH_____o_____o_____>

5

1260 1270 1280 1290 1300
acctgcaaat gagcagtctg aagtctgagg acacagcctt gttttactgt
tggacgttta ctcgtcagac ttcagactcc tgtgtcggaa caaatgaca
Y L Q M S S L K S E D T A L F Y C>

_____o_____o_VH_____o_____o_____>

10

1310 1320 1330 1340 1350
gcaagacacc gctcgggcta cttttctatg gactactggg gtcaaggaac
cgttctgtgg cgagcccgat gaaaagatac ctgatgaccc cagttccttg
A R H R S G Y F S M D Y W G Q G T>

15

_____o_____o_VH_____o_____o_____>

1360 1370 1380 1390 1400
ctcagtcacc gtctcctcat gccaccgtg cccagcacct gaactcctag
gagtcagtgg cagaggagta cgggtggcac gggtcgtgga cttgaggatc
S V T V S S>

20

_____VH_____>

A P E L L>

_____CH2_____>

C P P C P>

25

_s_HINGE_s_>

1410 1420 1430 1440 1450
ggggaccgtc agtcttcctc ttccccccaa aacccaagga caccctcatg
cccctggcag tcagaaggag aaggggggtt ttgggttcct gtgggagtac
G G P S V F L F P P K P K D T L M>

30

_____P_____P_____CH2_____P_____P_____>

1460 1470 1480 1490 1500
atctcccga cccctgaggt cacatgcgtg gtgggtggacg tgagccacga
tagagggcct ggggactcca gtgtacgcac caccacctgc actcgggtgct

35

- - 116 - -

I S R T P E V T C V V V D V S H E>
_____P_____P_____CH2_____P_____P_____>

5
1510 1520 1530 1540 1550
agaccctgag gtcaagttca actggtacgt ggacggcgtg gaggtgcata
tctgggactc cagttcaagt tgaccatgca cctgccgcac ctccacgtat
D P E V K F N W Y V D G V E V H>
_____P_____P_____CH2_____P_____P_____>

10
1560 1570 1580 1590 1600
atgccaagac aaagccgcgg gaggagcagt acaacagcac gtaccgggtg
tacggttctg ttctggcgcc ctctcgtca tgttgctcgtg catggcccac
N A K T K P R E E Q Y N S T Y R V>
_____P_____P_____CH2_____P_____P_____>

15
1610 1620 1630 1640 1650
gtcagcgtcc tcaccgtcct gcaccaggac tggctgaatg gcaaggagta
cagtcgcagg agtggcagga cgtggctcctg accgacttac cgttcctcat
V S V L T V L H Q D W L N G K E Y>
_____P_____P_____CH2_____P_____P_____>

20
1660 1670 1680 1690 1700
caagtgaag gtctccaaca aagccctccc agcccccatc gagtaactcg
gttcacgttc cagaggttgt ttcgggaggg tcgggggtag ctcatcgagc
* L>
_____>

25
K C K V S N K A L P A P I E>
_____P_____CH2_____P_____P_____>

30
ag
tc
E>
_____>

35

- - 117 - -

[SEQ. ID. NO.: 12] - Fiber Construct A1 C242

5 Sequence Range: 1 to 1111

```
>EcoR1
|
|      10      20      30      40      50
10  gaattcgatg aagcgcgcaa gaccgtctga agataccttc aaccccgtgt
    cttaagctac ttcgcgcggtt ctggcagact tctatggaag ttggggcaca
        M>
        ____>
        M  K  R  A  R  P  S  E  D  T  F  N  P  V>
15  ____o____o____TAIL____o____>

        60      70      80      90      100
    atccatatga cacggaaacc ggtcctccaa ctgtgccttt tcttactcct
    taggtatact gtgcctttgg ccaggagggtt gacacggaaa agaatgagga
20  Y  P  Y  D  T  E  T  G  P  P  T  V  P  F  L  T  P>
    ____o____o____TAIL____o____o____>

        110     120     130     140     150
    ccctttgtat cccccaatgg gtttcaagag agtccccctg ggggtactctc
25  gggaaacata gggggttacc caaagttctc tcaggggggac cccatgagag
                                           L  S>
                                           ____>

    P  F  V  S  P  N  G  F  Q  E  S  P  P  G  V>
    ____o____TAIL____o____o____>
30

        160     170     180     190     200
    tttgcgccta tccgaacctc tagttacctc caatggcatg cctgacgtag
    aaacgcggat aggcttggag atcaatggag gttaccgtac ggactgcac
    L  R  L  S  E  P  L  V  T  S  N  G  M>
35  ____m____REPEAT 1____m____>
```

- - 118 - -

P D V>

5

210 220 230 240 250
caagcttacg acaacaggta gaagccttgc aagggcaggt acaacactta
gttcgaatgc tgttgtccat cttcggaacg ttcccgtcca tgttgtgaat
A S L R Q Q V E A L Q G Q V Q H L>
LUNG SURFACTANT PROTEIN D TRIM_p>

10

260 270 280 290 300
caggcggcat ttagccaata caaaaaggta gagttgtttc caaacggagc
gtccgccgta aatcggttat gtttttccat ctcaacaaag gtttgcctcg
A>

15

Q A A F S Q Y K K V E L F P N G>
LUNG SURFACTANT PROTEIN D TRIM_p>

>Cla1

20

310 320 330 340 350
caagaagctg aacgacgccc agggccccc aa gagcgaccca tcgatcgata
gttcttcgac ttgctgcggg tccggggggt ctcgctgggt agctagctat
K K L N D A Q A P K S D>

25

b SPA 48-60 b>

P S I>

c>

D>

30

360 370 380 390 400
ttgtgatgac tcaggctgca ccctctgtac ctgtcactcc tggagagtca
aacactactg agtccgacgt gggagacatg gacagtgagg acctctcagt
I V M T Q A A P S V P V T P G E S>
e VK PHARMACIA e>

35

- - 119 - -

410 420 430 440 450
gtatccatct cctgcaggtc tagtaagagt ctctgcata gtaatggcaa
cataggtaga ggacgtccag atcatttctca gaggacgtat cattaccgtt
V S I S C R S S K S L L H S N G N>
5 _____e_____VK PHARMACIA_____e_____>
S L L H S N G N>
_____g_____CDR1_____>

460 470 480 490 500
cacttacttg tattgggttc tgcagaggcc aggccagtct cctcagctcc
gtgaatgaac ataaccaagg acgtctccgg tccggtcaga ggagtcgagg
T Y L Y W F L Q R P G Q S P Q L>
_____e_____VK PHARMACIA_____e_____>
T Y L Y>
15 _____g_____>

510 520 530 540 550
tgatatatcg gatgtccaac ctgtgtctcag gagtcccaga caggttcagt
actatatagc ctacagggttg gaacagagtc ctcaggggtct gtccaagtca
L I Y R M S N L V S G V P D R F S>
_____e_____VK PHARMACIA_____e_____>
R M S N L V S>
_____f_____CDR2_____>

560 570 580 590 600
ggcagtgggt caggaactgc tttcacactg agaatcagta gagtggaggc
ccgtcaccca gtccttgacg aaagtgtgac tcttagtcat ctcacctccg
G S G S G T A F T L R I S R V E A>
_____e_____VK PHARMACIA_____e_____>

610 620 630 640 650
tgaggatgtg ggtgtttatt actgtctgca acatctagag tatccgttca
actcctacac ccacaaataa tgacagacgt tgtagatctc ataggcaagt
L Q H L E Y P F>
35 _____d_____CDR3_d_____>

- - 120 - -

E D V G V Y Y C L Q H L E Y P F>

_____e_____VK PHARMACIA_____e_____>

5

660 670 680 690 700
 cgttcgggtcc tgggaccaag ctggagctga aacggccccc ggactttgtt
 gcaagccagg accctgggttc gacctcgact ttgccggggg cctgaaacaa
 T>

____>

10

T F G P G T K L E L K R>

_____e_____VK PHARMACIA_____e_____>

P P D F V>

_____825_____>

15

710 720 730 740 750
 cccccggccg ctagtttccc tgatcactcc cctcgtggcc aggtccagtt
 gggggccggc gatcaaaggg actagtgggg ggagcaccgg tccaggtcaa

Q V Q L>

____i_____>

20

P P A A S F P D H S P R G>

_____l_____825_____l_____>

25

760 770 780 790 800
 ggtgcagtct ggacctgagc tgaagaagcc tggagagaca gtcaagatct
 ccacgtcaga cctggactcg acttcttcgg acctctctgt cagttctaga
 V Q S G P E L K K P G E T V K I>

_____i_____VH PHARMACIA [SPLIT]_____i_____>

30

810 820 830 840 850
 cctgcaaggc ttctgattat accttcacat actatggaat gaactgggtg
 ggacgttccg aagactaata tggaagtgtg tgatacctta cttgaccac
 S C K A S D Y T F T Y Y G M N W V>

_____i_____VH PHARMACIA [SPLIT]_____i_____>

Y Y G M N>

____k_____CDR1____k_____>

35

860 870 880 890 900

- - 121 - -

aagcaggctc cgaggaaaggg tttaaagtgg atgggctgga tagacaccac
ttcgtccgag gccctttccc aaatttcacc tacccgacct atctgtggtg
K Q A P G K G L K W M G W I D T T>
_____i_____VH PHARMACIA [SPLIT]_____i_____>
5 W I D T T>
_____CDR2 [SP_____>
910 920 930 940 950
cactggagag ccaacatatg ctgaagattt taagggacgg attgccttct
10 gtgacctctc ggttggtatac gacttctaaa attccctgcc taacggaaga
T G E P T Y A E D F K G R I A F>
_____i_____VH PHARMACIA [SPLIT]_____i_____>
T G E P T Y A E D F K G>
_____j_____CDR2 [SPLIT]_____j_____>
15 960 970 980 990 1000
ctttggagac ctctgccagc actgcctatt tgcagatcaa aaacctcaaa
gaaacctctg gagacgggtcg tgacggataa acgtctagtt tttggagttt
S L E T S A S T A Y L Q I K N L K>
20 _____i_____VH PHARMACIA [SPLIT]_____i_____>
1010 1020 1030 1040 1050
aatgaggaca cggctacata tttctgtgca agacgggggc cttacaactg
ttactcctgt gccgatgtat aaagacacgt tctgcccccg gaatgttgac
25 R G P Y N W>
_____CDR3_____>
N E D T A T Y F C A R R G P Y N W>
_____i_____VH PHARMACIA [SPLIT]_____i_____>
1060 1070 1080 1090 1100
30 gtactttgat gtctggggcc aagggaccac ggtcaccgtc tctcactcg
catgaaacta cagaccccgg ttccctggtg ccagtggcag aggagtgagc
L>
_____>
Y F D V>
35 _____h_____>

- - 122 - -

Y F D V W G Q G T T V T V S S>
_____i____VH PHARMACIA [SPLIT]_____i____>

>Xho1

5

|

|1110

attaactcga g

taattgagct c

D * L E>

10

_____a_>

[SEQ. ID. NO.: 13] - Fiber Construct A7 C242

15

Sequence Range: 1 to 1402

>EcoR1

|

20

|

10

20

30

40

50

gaattcgatg aagcgcgcaa gaccgtctga agataccttc aaccccggtg
cttaagctac ttgcgcgctt ctggcagact tctatggaag ttggggcaca

M>

____>

25

M K R A R P S E D T F N P V>

____v____v____TAIL____v____>

60

70

80

90

100

30

atccatatga cacggaaacc ggtcctccaa ctgtgccttt tcttactcct
taggtatact gtgccttttg ccaggagggt gacacggaaa agaagtagga
Y P Y D T E T G P P T V P F L T P>

____v____v____TAIL____v____>

110

120

130

140

150

35

ccctttgtat cccccaatgg gtttcaagag agtccccctg gggtagctctc

- - 123 - -

gggaaacata ggggggttacc caaagttctc tcagggggac cccatgagag
L S>

P F V S P N G F Q E S P P G V>
5 _____v_____TAIL_____v_____v_____>
160 170 180 190 200
tttgcgcta tccgaacctc tagttacctc caatggcatg cttgcgctca
aaacgcggat aggcttggag atcaatggag gttaccgtac gaacgcgagt
10 L A L>

L R L S E P L V T S N G M>
_____t_____REPEAT 1_____t_____>
15 210 220 230 240 250
aatgggcaa cggcctctct ctggacgagg ccggcaacct tacctcccaa
tttaccggtt gccggagaga gacctgctcc ggccgttgga atggagggtt
L T S Q>
____r_____>
20 K M G N G L S L D E A G N>
_____s_____REPEAT 2_____s_____>
260 270 280 290 300
aatgtaacca ctgtgagccc acctctcaaa aaaaccaagt caaacataaa
25 ttacattggt gacactcggg tggagagttt ttttggttca gtttgtattt
I N>

N V T T V S P P L K K T K S N>
_____r_____REPEAT 3_____r_____r_____>
30 310 320 330 340 350
cctggaaata tctgcacccc tcacagttac ctgagaagcc ctaactgtgg
ggacctttat agacgtgggg agtgtcaatg gagtcttcgg gattgacacc
L T V>
35 _____>

- - 124 - -

L E I S A P L T V T S E A>
_____q_____REPEAT 4_____q_____>

5 360 370 380 390 400
ctgccgccgc acctctaata gtcgcgggca acacactcac catgcaatca
gacggcgggcg tggagattac cagcgcccgt tgtgtgagtg gtacgttagt
L T M Q S>
_____REPEAT 6_____>

10 A A A A P L M V A G N T>
_____P_____REPEAT 5_____P_____>

410 420 430 440 450
cagggccccgc taaccgtgca cgactccaaa cttagcattg ccaccaagg
gtccggggcg attggcacgt gctgagggtt gaatcgtaac ggtgggttcc
L S I A T Q G>
_____REPEAT 7_____>

Q A P L T V H D S K>
_____REPEAT 6_____o_____>

20 460 470 480 490 500
accctcaca gtgtcagaag gaaagctagc ccctgacgta gcaagcttac
tggggagtggt cacagtcttc ctttcgatcg gggactgcat cgttcgaatg
L A>
_____m_____>

25 P L T V S E G K>
_____REPEAT 7_____n_____>

P D V A S L>
_____LUNG SURFACTA_____>

30 510 520 530 540 550
gacaacagggt agaagccttg caagggcagg tacaacactt acaggcggca
ctgttggtcca tcttcggaac gttcccgtcc atgttggtgaa tgtccgccgt
R Q Q V E A L Q G Q V Q H L Q A A>
_____LUNG SURFACTANT PROTEIN D TRIM_____w_____>

35 560 570 580 590 600

- - 125 - -

tttagccaat acaaaaaggt agagttgttt ccaaacggag ccaagaagct
aaatcgggta tgtttttcca tctcaacaaa ggtttgccctc ggttcttcga
A K K L>
_b_____>
5 F S Q Y K K V E L F P N G>
_____LUNG SURFACTANT PROTEIN D TRIM_____>

>Cla1
|
10 610 620 630 | 640 650
gaacgacgcc caggccccca agagcgaccc atcgatcgat attgtgatga
cttgctgcgg gtccgggggt tctcgctggg tagctagcta taacactact
N D A Q A P K S D>
_____SPA 48-60_b_____>
15 P S I>
_c_____>
D I V M>
_e_____>
20 660 670 680 690 700
ctcaggctgc accctctgta cctgtcactc ctggagagtc agtatccatc
gagtcgcgacg tgggagacat ggacagtgag gacctctcag tcataggtag
T Q A A P S V P V T P G E S V S I>
_____e_____VK PHARMACIA_____e_____>
25 710 720 730 740 750
tcctgcaggt ctagtaagag tctcctgcat agtaatggca acacttactt
aggacgtcca gatcattctc agaggacgta tcattaccgt tgtgaatgaa
S C R S S K S L L H S N G N T Y L>
30 _____e_____VK PHARMACIA_____e_____>
S L L H S N G N T Y L>
_g_____g_CDR1_____g_____>
35 760 770 780 790 800
gtattgggtc ctgcagaggc caggccagtc tcctcagctc ctgatataac

- - 126 - -

cataaccaag gacgtctccg gtccgggtcag aggagtcgag gactatatag
Y W F L Q R P G Q S P Q L L I Y>
_____e_____VK PHARMACIA_____e_____>
5 Y>
____>

810 820 830 840 850
ggatgtccaa ccttgtctca ggagtcccag acaggttcag tggcagtggg
10 cctacagggtt ggaacagagt cctcagggtc tgtccaagtc accgtcaccc
R M S N L V S G V P D R F S G S G>
_____e_____VK PHARMACIA_____e_____>
R M S N L V S>
_____CDR2_____>
15

860 870 880 890 900
tcaggaactg ctttcacact gagaatcagt agagtggagg ctgaggatgt
agtccttgac gaaagtgtga ctcttagtca tctcacctcc gactcctaca
20 S G T A F T L R I S R V E A E D V>
_____e_____VK PHARMACIA_____e_____>

910 920 930 940 950
gggtgtttat tactgtctgc aacatctaga gtatccgttc acgttcggtc
cccacaaata atgacagacg ttgtagatct cataggcaag tgcaagccag
25 L Q H L E Y P F T>
_____d_____CDR3_____d_____>
G V Y Y C L Q H L E Y P F T F G>
_____e_____VK PHARMACIA_____e_____>
30

960 970 980 990 1000
ctgggaccaa gctggagctg aaacggcccc cggactttgt tccccggcc
gaccctgggt cgacctcgac ttgcccgggg gcctgaaaca agggggccgg
P G T K L E L K R>
35 _____VK PHARMACIA_____e_____>
P P D F V P P A>

- - 127 - -

_____1_____825_1_____>

1010 1020 1030 1040 1050
 gctagtttcc ctgatcactc ccctcgtggc caggtccagt tgggtgcagtc
 5 cgatcaaagg gactagtgag gggagcaccg gtccagggtca accacgtcag
 Q V Q L V Q S>
 ____VH PHARMACIA [____>

A S F P D H S P R G>
 _____1_____825_____1_____>

10

1060 1070 1080 1090 1100
 tggacctgag ctgaagaagc ctggagagac agtcaagatc tcctgcaagg
 acctggactc gacttcttcg gacctctctg tcagttctag aggacgttcc
 G P E L K K P G E T V K I S C K>
 15 _____i_____VH PHARMACIA [SPLIT]_____i_____>

1110 1120 1130 1140 1150
 cttctgatta taccttcaca tactatggaa tgaactgggt gaagcaggct
 gaagactaat atggaagtgt atgatacctt acttgaccca cttcgtccga
 20 A S D Y T F T Y Y G M N W V K Q A>
 _____i_____VH PHARMACIA [SPLIT]_____i_____>
 Y Y G M N>
 _____CDR1_k_____>

25

1160 1170 1180 1190 1200
 ccgggaaagg gtttaaagtg gatgggctgg atagacacca ccactggaga
 ggccctttcc caaatttcac ctacccgacc tatctgtggt ggtgacctct
 P G K G L K W M G W I D T T T G E>
 _____i_____VH PHARMACIA [SPLIT]_____i_____>
 30 W I D T T T G E>
 ____j____CDR2 [SPLIT]_____>

1210 1220 1230 1240 1250
 gccaacatat gctgaagatt ttaagggacg gattgccttc tctttggaga
 35 cggttgtata cgacttctaa aattccctgc ctaacggaag agaaacctct

- - 128 - -

P T Y A E D F K G R I A F S L E>
_____i_____VH PHARMACIA [SPLIT]_____i_____>
P T Y A E D F K G>
_____CDR2 [SPLIT]_j_____>
5

1260 1270 1280 1290 1300
cctctgccag cactgcctat ttgcagatca aaaacctcaa aaatgaggac
ggagacggtc gtgacggata aacgtctagt ttttgagatt tttactcctg
T S A S T A Y L Q I K N L K N E D>
10 _____i_____VH PHARMACIA [SPLIT]_____i_____>

1310 1320 1330 1340 1350
acggctacat atttctgtgc aagacggggg ccttacaact ggtactttga
tgccgatgta taaagacacg ttctgcccc ggaatggtga ccatgaaact
15 R G P Y N W Y F D>
_____h_____CDR3_h_____>
T A T Y F C A R R G P Y N W Y F D>
_____i_____VH PHARMACIA [SPLIT]_____i_____>

20 >Xho1
|
1360 1370 1380 1390 1400
tgtctggggc caagggacca cggtcaccgt ctctcactc gattaactcg
acagaccccc gttccctggt gccagtggca gaggagttag ctaattgagc
25 L D * L>
_____a_____>

V>

V W G Q G T T V T V S S>
30 _____VH PHARMACIA [SPLIT]_____i_____>

ag
tc
E>
35 _____>

- - 129 - -

[SEQ. ID. NO.: 14] -

5 Sequence Range: 1 to 174 - Affibody ZIgG

```

          10          20          30          40          50
gtagacaaca aattcaacaa agaacaacaa aacgcgttct atgagatctt
catctgttgt ttaagttggt tcttggtggt ttgcgcaaga tactctagaa
10  V D N K F N K E Q Q N A F Y E I L>
      _____a_____a_Z-DOM?N__a_____a_____>

          60          70          80          90          100
acatttacct aacttaaacg aagaacaacg aaacgccttc atccaaagtt
15  tgtaaagtga ttgaatttgc ttcttggtgc tttgcggaag taggtttcaa
      H L P N L N E E Q R N A F I Q S>
      _____a_____a_Z-DOM?N__a_____a_____>

          110          120          130          140          150
taaaagatga cccaagccaa agcgctaact tgctagcaga agctaaaaag
20  attttctact gggttcgggt tgcgcattga acgatcgtct tcgatttttc
      L K D D . P S Q S A N L L A E A K K>
      _____a_____a_Z-DOM?N__a_____a_____>

          160          170
ctaaatgatg ctcaggcgcc gaaa
25  gatttactac gagtccgcgg cttt
      L N D A Q A P K>
      _____Z-DOM?N_____a_____>
30
```

[SEQ. ID. NO.: 15] - Affibody ZIgA

35 Sequence Range: 1 to 174

RECTIFIED SHEET (RULE 91)

- - 130 - -

10 20 30 40 50
gtagacaaca aattcaacaa agaaacaata caagcgagtc aagagatcag
catctgttgt ttaagttggt tctttgttat gttcgctcag ttctctagtc
V D N K F N K E T I Q A S Q E I R>
5 _____a_____AFFI IGA _a_____a_____>
60 70 80 90 100
actattacct aacttaaacg gtagacaaaa gcttgccttc atccacagtt
tgataatgga ttgaatttgc catctgtttt cgaacggaag taggtgtcaa
10 L L P N L N G R Q K L A F I H S>
_____a_____AFFI IGA _a_____a_____>
110 120 130 140 150
tacttgatga cccaagccaa agcgctaact tgctagcaga agctaaaaag
15 atgaactact gggttcgggt tcgcgattga acgatcgtct tcgatttttc
L L D D P S Q S A N L L A E A K K>
_____a_____AFFI IGA _a_____a_____>
160 170
20 ctaaattgatg ctcaggcgcc gaaa
gatttactac gagtccgcgg cttt
L N D A Q A P K>
_____AFFI IGA ____a____>

25

[SEQ. ID. NO.: 16] - Recombinant Fiber A7 ZIgG

Sequence Range: 1 to 820

30

35

>EcoR1

|

|

10 20 30 40 50
gaattcgatg aagcgcgcaa gaccgtctga agataccttc aaccccggtg
cttaagctac ttcgcgcggt ctggcagact tctatggaag ttggggcaca

- - 131 - -

M>
____>
M K R A R P S E D T F N P V>
____j____j____TAIL____j____>

5
60 70 80 90 100
atccatatga cacggaaacc ggtcctccaa ctgtgccttt tcttactcct
taggtatact gtgcctttgg ccaggagggtt gacacggaaa agaagtagga
Y P Y D T E T G P P T V P F L T P>
10 _____j____j____TAIL____j____j____>

110 120 130 140 150
ccctttgtat cccccaatgg gtttcaagag agtccccctg gggtagctct
gggaaacata ggggggtacc caaagttctc tcaggggggac cccatgagag
15 _____L S>
____>
P F V S P N G F Q E S P P G V>
____j____TAIL____j____j____>

20 160 170 180 190 200
tttgcgccta tccgaacctc tagttacctc caatggcatg cttgcgctca
aaacgcggat aggcttggag atcaatggag gttaccgtac gaacgcgagt
L A L>
____>

25 L R L S E P L V T S N G M>
____h____REPEAT 1____h____>

30 210 220 230 240 250
aaatgggcaa cggcctctct ctggacgagg ccggcaacct tacctcccaa
tttaccggtt gccggagaga gacctgctcc ggccgttgga atggagggtt
L T S Q>
____f____>

K M G N G L S L D E A G N>
____g____REPEAT 2____g____>

35

- - 132 - -

260 270 280 290 300
aatgtaacca ctgtgagccc acctctcaaa aaaaccaagt caaacataaa
ttacattggt gacactcggg tggagagttt ttttggttca gtttgtattt
I N>
5 N V T T V S P P L K K T K S N>
f REPEAT 3 f f >
310 320 330 340 350
cctggaaata tctgcacccc tcacagttac ctcagaagcc ctaactgtgg
ggacctttat agacgtgggg agtgtcaatg gagtcttcgg gattgacacc
L T V>
10 L E I S A P L T V T S E A>
e REPEAT 4 e >
15 360 370 380 390 400
ctgccgccgc acctctaattg gtcgcgggca acacactcac catgcaatca
gacggcgggcg tggagattac cagcgcccgt tgtgtgagtg gtacgttagt
L T M Q S>
20 REPEAT 6 >
A A A A P L M V A G N T>
d REPEAT 5 d >
25 410 420 430 440 450
caggccccgc taaccgtgca cgactccaaa cttagcattg ccacccaagg
gtccggggcg attggcacgt gctgaggttt gaatcgtaac ggtgggttcc
L S I A T Q G>
30 REPEAT 7 >
Q A P L T V H D S K>
REPEAT 6 c >
35 460 470 480 490 500
accctcaca gtgtcagaag gaaagctagc ccctgacgta gcaagcttac
tggggagtggt cacagtcttc ctttcgatcg gggactgcat cgttcgaatg

- - 133 - -

L A>
____a_>
P L T V S E G K>
____REPEAT 7____b____>
5 P D V A S L>
____LUNG SURFACTA____>
510 520 530 540 550
gacaacaggt agaagccttg caagggcagg tacaacactt acaggcggca
10 ctgttgcca tcttcggaac gttcccgtcc atgttgatgaa tgtccgccgt
R Q Q V E A L Q G Q V Q H L Q A A>
____LUNG SURFACTANT PROTEIN D TRIM____k____>
560 570 580 590 600
15 tttagccaat acaaaaaggt agagttgttt ccaaaccggag ccaagaagct
aaatcggtta tgtttttcca tctcaacaaa ggtttgccctc gggttcttcga
F S Q Y K K V E L F P N G>
____LUNG SURFACTANT PROTEIN D TRIM____>
A K K L>
20 ____l____>
>Cla1
|
610 620 630 | 640 650
25 gaacgacgcc caggccccca agagcgaccc atcgatcgta gacaacaaat
cttgctgcgg gtccgggggt tctcgctggg tagctagcat ctgttggtta
N D A Q A P K S D>
____SPA 48-60____l____>
P S I>
30 ____m____>
V D N K>
____o____>
660 670 680 690 700
35 tcaacaaaga acaacaaaac gcgttctatg agatcttaca ttacctaac

- - 134 - -

agttgtttct tgttgttttg cgcaagatac tctagaatgt aaatggattg
F N K E Q Q N A F Y E I L H L P N>
_____o_____o_Z-DOM?N_____o_____o_____>

5 710 720 730 740 750
ttaaacgaag aacaacgaaa cgccttcatac caaagtttaa aagatgaccc
aatttgcttc ttgttgcttt gcggaagtag gtttcaaatt ttctactggg
L N E E Q R N A F I Q S L K D D P>
_____o_____o_Z-DOM?N_____o_____o_____>

10 760 770 780 790 800
aagccaaagc gctaacttgc tagcagaagc taaaaagcta aatgatgctc
ttcgggtttcg cgattgaacg atcgtcttcg atttttcgat ttactacgag
S Q S A N L L A E A K K L N D A>
_____o_____o_Z-DOM?N_____o_____o_____>

15 >Xho1
 |
 810 | 820
aggcgccgaa ataactcgag
20 tccgcggctt tattgagctc
 *>
 ____>
 Q A P K>
 _____o_____>

25

[SEQ. ID. NO.: 17] - Recombinant Fiber A7 ZIgA

30

Sequence Range: 1 to 820

>EcoR1
|
35 | 10 20 30 40 50

- - 135 - -

gaattcgatg aagcgcgcaa gaccgtctga agataccttc aaccccgtgt
cttaagctac ttcgcgcggtt ctggcagact tctatggaag ttggggcaca

M>

____>

5

M K R A R P S E D T F N P V>

____j____j____TAIL____j____>

60

70

80

90

100

10

atccaatgatga cacggaaacc ggtcctccaa ctgtgccttt tcttactcct
taggtatact gtgcctttgg ccaggagggtt gacacggaaa agaattgagga
Y P Y D T E T G P P T V P F L T P>

____j____j____TAIL____j____>

110

120

130

140

150

15

ccctttgtat cccccaatgg gtttcaagag agtccccctg ggggtactctc
gggaaacata ggggggttacc caaagttctc tcaggggggac cccatgagag

L S>

____>

P F V S P N G F Q E S P P G V>

20

____j____TAIL____j____>

160

170

180

190

200

25

tttgcgccta tccgaacctc tagttacctc caatggcatg cttgcgctca
aaacgcggat aggcttggag atcaatggag gttaccgtac gaacgcgagt

L A L>

____>

L R L S E P L V T S N G M>

____h____REPEAT 1____h____>

30

210

220

230

240

250

aaatgggcaa cggcctctct ctggacgagg ccggcaacct tacctcccaa
tttaccggtt gccggagaga gacctgctcc ggccggttga atggagggtt

L T S Q>

____f____>

35

K M G N G L S L D E A G N>

- - 136 - -

BNSDOCID: <WO_____0208263A2 | >

- - 137 - -

accctcaca gtgtcagaag gaaagctagc ccctgacgta gcaagcttac
tggggagtggt cacagtcttc ctttcgatcg gggactgcat cgttcgaatg

L A>

_____a_>

5

P L T V S E G K>

_____REPEAT 7_____b_____>

P D V A S L>

_____LUNG SURFACTANT_____>

10

510 520 530 540 550
gacaacaggt agaagccttg caagggcagg tacaacactt acaggcggca
ctgttggtcca tcttcggaac gttcccgtcc atgttgtgaa tgtccgccgt
R Q Q V E A L Q G Q V Q H L Q A A>

_____LUNG SURFACTANT PROTEIN D TRIM_____k_____>

15

560 570 580 590 600
tttagccaat acaaaaaggt agagttgttt ccaaacggag ccaagaagct
aaatcgggtta tgtttttcca tctcaacaaa ggtttgcctc ggttcttcga
F S Q Y K K V E L F P N G>

20

_____LUNG SURFACTANT PROTEIN D TRIM_____>

A K K L>

_____l_____>

>ClaI

25

610 620 630 | 640 650
gaacgacgcc caggccccca agagcgaccc atcgatcgta gacaacaaat
cttgctgcgg gtccgggggt tctcgctggg tagctagcat ctgttggtta
N D A Q A P K S D>

30

_____SPA 48-60_____l_____>

P S I>

_____m_____>

V D N K>

_____o_____>

35

- - 138 - -

5 660 670 680 690 700
tcaacaaaga aacaatacaa gcgagtcaag agatcagact attacctaac
agttgtttct ttgttatggt cgctcagttc tctagtctga taatggattg
F N K E T I Q A S Q E I R L L P N>
_____o_____AFFI IGA _o_____o_____>

10 710 720 730 740 750
ttaaacggta gacaaaagct tgccttcatt cacagtttac ttgatgaccc
aatttgccat ctgttttcga acggaagtag gtgtcaaagt aactactggg
L N G R Q K L A F I H S L L D D P>
_____o_____AFFI IGA _o_____o_____>

15 760 770 780 790 800
aagccaaagc gctaacttgc tagcagaagc taaaaagcta aatgatgctc
ttcgggtttcg cgattgaacg atcgtcttcg atttttcgat ttactacgag
S Q S A N L L A E A K K L N D A>
_____o_____AFFI IGA _o_____o_____>

20 >Xho1
 |
 810 | 820
aggcgccgaa ataactcgag
tccgcggctt tattgagctc
 *>
25 ____>
 Q A P K>
 ____o____>

30

[SEQ. ID. NO.: 18] - Recombinant Fiber A7 ZIgG/ZIgA

Sequence Range: 1 to 1003

35

- - 139 - -

>EcoR1

5
|
| 10 20 30 40 50
gaattcgatg aagcgcgcaa gaccgtctga agataccttc aaccccgtgt
cttaagctac ttcgcgcggtt ctggcagact tctatggaag ttgggggcaca
M>
____>
M K R A R P S E D T F N P V>
____j____j____TAIL____j____>
10
60 70 80 90 100
atccatatga cacggaaacc ggtcctccaa ctgtgccttt tcttactcct
taggtatact gtgccttttg ccaggagggtt gacacggaaa agaatgagga
Y P Y D T E T G P P T V P F L T P>
15
____j____j____TAIL____j____>
110 120 130 140 150
ccctttgtat cccccaatgg gtttcaagag agtccccctg ggggtactctc
gggaaacata ggggggttacc caaagttctc tcaggggggac cccatgagag
20
L S>
____>
P F V S P N G F Q E S P P G V>
____j____TAIL____j____j____>
25
160 170 180 190 200
tttgcgccta tccgaacctc tagttacctc caatggcatg cttgcgctca
aaacgcggat aggcttggag atcaatggag gttaccgtac gaacgcgagt
L A L>
____>
30
L R L S E P L V T S N G M>
____h____REPEAT 1____h____>
210 220 230 240 250
aaatgggcaa cggcctctct ctggacgagg cgggcaacct tacctcccaa
35
tttaccggtt gccggagaga gacctgctcc ggccggttga atggagggtt

- - 140 - -

5
L T S Q>
____f____>
K M G N G L S L D E A G N>
____g____REPEAT 2____g____>

260 270 280 290 300
aatgtaacca ctgtgagccc acctctcaaa aaaaccaagt caaacataaa
ttacattggt gacactcggg tggagagttt ttttggttca gtttgatattt
I N>
____>

10
N V T T V S P P L K K T K S N>
____f____REPEAT 3____f____f____>

310 320 330 340 350
cctggaaata tctgcacccc tcacagttac ctcagaagcc ctaactgtgg
ggacctttat agacgtgggg agtgtcaatg gagtcttcgg gattgacacc
L T V>
____>

20
L E I S A P L T V T S E A>
____e____REPEAT 4____e____>

360 370 380 390 400
ctgccgccgc acctctaata gtcgcgggca acacactcac catgcaatca
gacggcggcg tggagattac cagcgcccgt tgtgtgagtg gtacgttagt
L T M Q S>
____REPEAT 6____>

25
A A A A P L M V A G N T>
____d____REPEAT 5____d____>

30
410 420 430 440 450
caggccccgc taaccgtgca cgactccaaa cttagcattg ccacccaagg
gtccggggcg attggcacgt gctgagggtt gaatcgtaac ggtggggttcc
L S I A T Q G>
____REPEAT 7____>

35
Q A P L T V H D S K>

- - 141 - -

____REPEAT 6_c____>

460 470 480 490 500
acccctcaca gtgtcagaag gaaagctagc ccctgacgta gcaagcttac
5 tgggggagtgt cacagtcttc ctttcgatcg gggactgcat cgttcgaatg
L A>
____a____>
P L T V S E G K>
____REPEAT 7_b____>

10 P D V A S L>
____LUNG SURFACTANT____>

510 520 530 540 550
gacaacaggt agaagccttg caagggcagg tacaacactt acaggcggca
15 ctgttggtcca tcttcggaac gttcccggtcc atgttggtgaa tgtccgccgt
R Q Q V E A L Q G Q V Q H L Q A A>
____LUNG SURFACTANT PROTEIN D TRIM_k____>

560 570 580 590 600
tttagccaat acaaaaaggt agagttgttt ccaaacggag ccaagaagct
20 aaatcggtta tgttttttcca tctcaacaaa ggtttgcttc ggtttcttga
F S Q Y K K V E L F P N G>
____LUNG SURFACTANT PROTEIN D TRIM____>

25 A K K L>
_l____>

>Cla1
|

610 620 630 | 640 650
30 gaacgacgcc caggccccca agagcgaccc atcgatcgta gacaacaaat
cttgctgcgg gtccgggggt tctcgctggg tagctagcat ctgttggtta
N D A Q A P K S D>
____SPA 48-60_l____>

35 P S I>
____m____>

- - 142 - -

V D N K>

____o____>

5

660 670 680 690 700
tcaacaaaga acaacaaaac gcgttctatg agatcttaca ttacctaac
agttgtttct tgttgttttg cgcaagatac tctagaatgt aaatggattg
F N K E Q Q N A F Y E I L H L P N>
____o____o_Z-DOM?N____o____o____>

10

710 720 730 740 750
ttaaacgaag aacaacgaaa cgccttcata caaagtttaa aagatgaccc
aatttgcttc ttgttgcttt gcggaagtag gtttcaaatt ttctactggg
L N E E Q R N A F I Q S L K D D P>
____o____o_Z-DOM?N____o____o____>

15

760 770 780 790 800
aagccaaagc gctaacttgc tagcagaagc taaaaagcta aatgatgctc
ttcggtttcg cgattgaacg atcgtcttcg atttttcgat ttactacgag
S Q S A N L L A E A K K L N D A>
____o____o_Z-DOM?N____o____o____>

20

810 820 830 840 850
aggcgccgaa aagctcgacc gtagacaaca aattcaacaa agaaacaata
tccgcggctt ttcgagctgg catctgttgt ttaagttggt tctttgttat
Q A P K>
____o____>

25

S S T>

____>

30

V D N K F N K E T I>

____AFFI IGA _q____>

35

860 870 880 890 900
caagcgagtc aagagatcag actattacct aacttaaagc gtagacaaaa
gttcgctcag ttctctagtc tgataatgga ttgaatttgc catctgtttt
Q A S Q E I R L L P N L N G R Q K>

- - 143 - -

_____q_____AFFI IGA _q_____q_____>

910 920 930 940 950

gcttgccttc atccacagtt tactttgatga cccaagccaa agcgctaact

5 cgaacggaag taggtgtcaa atgaactact gggttcgggtt tcgcgattga

L A F I H S L L D D P S Q S A N>

_____q_____AFFI IGA _q_____q_____>

'

>XhoI

10 |

960 970 980 990 1000

tgctagcaga agctaaaaag ctaaattgatg ctcaggcgcc gaaataactc

acgatcgtct tcgatttttc gatttactac gagtccgcgg ctttattgag

*>

15 _____>

L L A E A K K L N D A Q A P K>

_____q_____AFFI IGA _____q_____q_____>

gag

20 ctc

[SEQ. ID. NO.: 19] - Recombinant Fiber A7 ZIgG/ZIgG

25

Sequence Range: 1 to 1003

>EcoRI

30 |

| 10 20 30 40 50

gaattcgatg aagcgcgcaa gaccgtctga agataccttc aaccccgtgt

cttaagctac ttgcgcggtt ctggcagact tctatggaag ttgggggcaca

M>

35 _____>

- - 144 - -

M K R A R P S E D T F N P V>
____j____j____TAIL____j____>

5 60 70 80 90 100
atccatatga cacggaacc ggtcctccaa ctgtgccttt tcttactcct
taggtatact gtgcctttgg ccaggagggtt gacacggaaa agaattgagga
Y P Y D T E T G P P T V P F L T P>
____j____j____TAIL____j____>

10 110 120 130 140 150
ccctttgtat cccccaatgg gtttcaagag agtccccctg gggactctc
gggaaacata ggggggttacc caaagttctc tcagggggac cccatgagag
L S>
____>

15 P F V S P N G F Q E S P P G V>
____j____TAIL____j____>

20 160 170 180 190 200
tttgcgccta tccgaacctc tagttacctc caatggcatg cttgcgctca
aaacgcggat aggcttggag atcaatggag gttaccgtac gaacgcgagt
L A L>
____>

25 L R L S E P L V T S N G M>
____h____REPEAT 1____h____>

30 210 220 230 240 250
aaatgggcaa cggcctctct ctggacgagg ccggcaacct tacctcccaa
tttaccggtt gccggagaga gacctgctcc ggccggttga atggagggtt
L T S Q>
____f____>

35 K M G N G L S L D E A G N>
____g____REPEAT 2____g____>

260 270 280 290 300
aatgtaacca ctgtgagccc acctctcaaa aaaaccaagt caaacataaa

- - 145 - -

ttacattggt gacactcggg tggagagttt ttttggttca gtttgtatatt

I N>

>

N V T T V S P P L K K T K S N>

5

f REPEAT 3 f f >

310 320 330 340 350

cctggaaata tctgcacccc tcacagttac ctcagaagcc ctaactgtgg
ggacdtttat agacgtgggg agtgtcaatg gagtcttcgg gattgacacc

10

L T V>

>

L E I S A P L T V T S E A>

e REPEAT 4 e >

15

360 370 380 390 400

ctgccgccgc acctctaattg gtcgcgggca acacactcac catgcaatca
gacggcgggcg tggagattac cagcgcccgt tgtgtgagtg gtacgttagt

L T M Q S>

REPEAT 6 >

20

A A A A P L M V A G N T>

d REPEAT 5 d >

410 420 430 440 450

caggccccgc taaccgtgca cgactccaaa cttagcattg ccacccaagg
gtccggggcg attggcacgt gctgaggttt gaatcgtaac ggtgggttcc

25

L S I A T Q G>

REPEAT 7 >

Q A P L T V H D S K>

REPEAT 6 c >

30

460 470 480 490 500

accctcaca gtgtcagaag gaaagctagc ccctgacgta gcaagcttac
tggggagtgt cacagtcttc ctttcgatcg gggactgcat cgttcgaatg

L A>

35

a >

- - 146 - -

P L T V S E G K>
 _____REPEAT 7_____b_____>

P D V A S L>
 _____LUNG SURFACTA_____>

5

510 520 530 540 550
 gacaacaggt agaagccttg caagggcagg tacaacactt acagggcggca
 ctgttggtcca tcttcggaac gttcccgtcc atgttggtgaa tgtccgccgt
 R Q Q . V E A L Q G Q V Q H L Q A A>
 _____LUNG SURFACTANT PROTEIN D TRIM_____k_____>

10

560 570 580 590 600
 tttagccaat acaaaaaggt agagttgttt ccaaacggag ccaagaagct
 aaatcgggtta tgttttttcca tctcaacaaa ggtttgcctc ggtttcttcga
 F S Q Y K K V E L F P N G>
 _____LUNG SURFACTANT PROTEIN D TRIM_____>

15

A K K L>
 _____l_____>

20

>Cla1

610 620 630 | 640 650
 gaacgacgcc caggccccca agagcgaccc atcgatcgta gacaacaaat
 cttgctgcgg gtccgggggt tctcgctggg tagctagcat ctgttggttta
 N D A Q A P K S D>

25

_____SPA 48-60_____l_____>

P S I>
 _____m_____>

30

V D N K>
 _____o_____>

660 670 680 690 700
 tcaacaaaga acaacaaaac gcgttctatg agatcttaca tttacctaac
 agttgtttct tgttgttttg cgcaagatac tctagaatgt aaatggattg
 F N K E Q Q N A F Y E I L H L P N>

35

- - 147 - -

_____o_____o_Z-DOM?N_____o_____o_____>

710 720 730 740 750
ttaaacgaag aacaacgaaa cgccttcac caaagttaa aagatgaccc
5 aatttgcttc ttgttgcttt gcggaagtag gtttcaaatt ttctactggg
L N E E Q R N A F I Q S L K D D P>

_____o_____o_Z-DOM?N_____o_____o_____>

760 770 780 790 800
10 aagccaaagc gctaacttgc tagcagaagc taaaaagcta aatgatgctc
ttcggtttcg cgattgaacg atcgtcttcg atttttcgat ttactacgag
S Q S A N L L A E A K K L N D A>

_____o_____o_Z-DOM?N_____o_____o_____>

810 820 830 840 850
15 aggcgccgaa aagctcgacc gtagacaaca aattcaacaa agaacaacaa
tccgcggctt ttcgagctgg catctgttgt ttaagttggt tcttggttgt
Q A P K>

_____o_____>

20 V D N K F N K E Q Q>
_____p_Z-DOM?N_____p_____>

S S T>
_____>

860 870 880 890 900
25 aacgcgttct atgagatctt acatttacct aacttaaacg aagaacaacg
ttgcgcaaga tactctagaa tgtaaagga ttgaatttgc ttcttggtgc
N A F Y E I L H L P N L N E E Q R>

_____p_____p_Z-DOM?N_____p_____p_____>

30 910 920 930 940 950
aaacgccttc atccaaagtt taaaagatga cccaagccaa agcgctaact
tttgcggaag taggtttcaa attttctact gggttcgggt tcgcgattga
N A F I Q S L K D D P S Q S A N>

35 _____p_____p_Z-DOM?N_____p_____p_____>

- - 148 - -

5
10
15
20
25
30

960 970 980 990 1000

tgctagcaga agctaaaaag ctaaatagatg ctcaggcgcc gaaataactc
acgatcgtct tcgatttttc gatttactac gagtccgcgg ctttattgag

*>

L L A E A K K L N D A Q A P K>

_____P_____Z-DOM?N_____P_____P_____>

gag
ctc

SEQ I.D. No.: 47 - Anti β -galactosidase single chain Fv fragment

ATGGCCGAGGTGCAGCTGGTGGAGTCTGGGGGAAGCCTGGTCAAGCCTGGGGGG
TCCCTGAGACTCTCCTGTGCAGCCTCTGGATTACCTTCAGTAAGTATAGCATGAAC
TGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTGGATCTCATCCATTAGTGGTAG
TAGTAGATACATATACTACGCAGACTTCGTGAAGGGCCGATTACCATCTCCAGAGAC
AACGCCACGAACTCACTGTACCTGCAAATGAACAGCCTGAGAGCCGAGGACACGGCTG
TTTATTACTGTGTGAGATCCAGTATTACGATTTTTTGGTGGCGGTATGGACGTCTGG
GGCAGAGGCACCCTGGTCACCGTCTCCTCAGGTGGAGGCGGTTCAGGCGGAGGTGG
CAGCGGCGGTGGCGGATCGCAGTCTGTGCTGACTCAGCCTGCCTCCGTGTCTGGGT
CTCCTGGACAGTCGATCACCATCTCCTGCGCTGGAACCAGCAGTGACGTTGGTGGT
TATAACTATGTCTCCTGGTACCAACAACACCCAGGCAAAGCCCCCAAACATCATGAT
TTATGAGGACAGTAAGCGGCCCTCAGGGGTTTCTAATCGCTTCTCTGGCTCCAAGT
CTGGCAACACGGCCTCCCTGACAATCTCTGGGCTCCAGGCTGAGGACGAGGCTGAT
TATTACTGCAGCTCATATACAACCAGGAGCACTCGAGTTTTTCGGCGGAGGGACCAA
GCTGGCCGTCCTAGGTGCGGCCGCAGAACAAAACTCATCTCAGAAGAGGATCTGA
ATGGGGCCGCACATCACCATCATCACCAT

Claims:

1. A modified virus comprising one or more non-native polypeptides, which polypeptide comprises one or more
5 framework moieties each containing one or more binding moieties, which polypeptide is capable of being expressed in the cytoplasm and nucleus of a mammalian host cell in a conformation which is maintained in the absence of a ligand for said binding moieties, said conformation
10 allowing said binding moieties subsequently to bind with said ligand, and which polypeptide is capable of transport through the nuclear membrane, wherein said modified virus has an altered tropism conferred by said binding moieties.
- 15 2. The modified virus of claim 1 which is derived from a virus selected from the group consisting of adenoviruses, retroviruses, lentiviruses, adeno-associated viruses, Reoviridae, Picornaviridae, Parvoviridae, Papovaviridae and Caliciviridae.
- 20 3. The modified virus of claim 1 or claim 2 which is derived from human adenovirus
4. The modified virus of any one of claims 1 to 3 which
25 is derived from human adenovirus serotype 5.
5. The modified virus of any one of claims 1 to 4 wherein said non-native polypeptide replaces, is incorporated into, or forms a fusion protein with, a viral
30 protein component of the wild type virus.
6. The modified virus of any one of claims 1. to 5 wherein said non-native polypeptide and/or the viral component protein comprising the non-native polypeptide is
35 soluble in a cellular environment.

- - 150 - -

- 5 7. The modified virus of claim 6 wherein greater than 30% of the non-native polypeptide or the viral component protein comprising the non-native polypeptide is present in the soluble fraction of cell lysates of cells expressing the non-native polypeptide or viral component protein comprising the non-native polypeptide.
- 10 8. The modified virus of any one of claims 5 to 7 wherein said viral protein component is an adenoviral fiber protein.
- 15 9. The modified virus of claim 8 wherein said non-native polypeptide is incorporated into an adenoviral fiber protein such that the wild-type fiber knob or cell binding domain thereof is removed.
- 20 10. The modified virus of any one of claims 1 to 9 wherein said non-native polypeptide comprises one or more further elements that mimic the native structure or function of a viral component protein in which said non-native polypeptide is incorporated or which said non-native polypeptide replaces.
- 25 11. The modified virus of any one of claims 1 to 10 wherein said non-native polypeptide does not contain any di-sulphide bonds.
- 30 12. The modified virus of any one of claims 1 to 11 wherein the non-native polypeptide comprises one or more α -helical structures.
- 35 13. The modified virus of any one of claims 1 to 12 wherein said non-native polypeptide comprises a framework moiety derived from an antibody which is capable of productive folding in the cytoplasm and subsequent

- - 151 - -

transport into the cell nucleus.

14. The modified virus of claim 13 wherein said framework moiety comprises one or more binding moieties derived from
5 a CDR of a further antibody.

15. The modified virus of any one of claims 1 to 12 wherein said non-native polypeptide is or comprises a combinatorial protein or an affibody.
10

16. The modified virus of any one of claims 1 to 15 wherein said non-native polypeptide comprises one or more framework moieties derived from a bacterial receptor.

15 17. The modified virus of any one of claims 1 to 12 or 16 wherein said non-native polypeptide comprises one or more binding moieties which are present within one or more loops of a helical bundle and/or one or more of the loops connecting these bundles.

20 18. The modified virus of claim 16 wherein said non-native polypeptide comprises one or more framework moieties derived from the immunoglobulin binding Z-domain from staphylococcal protein A or an immunoglobulin binding
25 domain from streptococcal protein G.

19. The modified virus of any one of claims 1 to 18 wherein said non-native polypeptide comprises one or more framework moieties which comprise a non native
30 trimerisation motif.

20. The modified virus of claim 19 wherein said non-native polypeptide comprises one or more framework moieties which comprise the neck region peptide from human
35 lung surfactant protein D.

21. The modified virus of any one of claims 1 to 20 which comprises two or more different non-native polypeptides.
- 5 22. The modified virus of claim 21 which comprises a first non-native polypeptide which binds a target cell and a second non-native polypeptide which binds a production cell or permissive cell.
- 10 23. The modified virus of any one of claims 1 to 22 wherein said non-native polypeptide comprises a cleavage site positioned in a location that enables a binding moiety of the non-native polypeptide to be cleaved from the modified virus.
- 15 24. The modified virus of any one of claims 1 to 23 wherein said virus comprises a modified viral component comprising said non-native polypeptide and a corresponding unmodified viral component e.g. a wild-type fiber and a
- 20 25. The modified virus of any one of claims 1 to 24 wherein said non-native polypeptide comprises a binding moiety capable of binding to a cell specific ligand which may optionally be Prostate Specific Membrane Antigen, EGF receptor, Her-2/Neu, VEGF receptor, CD22, gp120, MHC/peptide complexes or membrane structures or surface molecules expressed or present on proliferating cells, tumor cells or virus infected cells.
- 30 26. The modified virus of any one of claims 1 to 25 which further comprises a site for insertion of one or more desired therapeutic genes or nucleic acid molecules.
- 35 27. The modified virus of claim 26 which comprises

- - 153 - -

transgenes encoding cytosine deaminase and/or uracil phosphoribosyl transferase either as separate genes or together as a bifunctional fusion gene.

5 28. The modified virus of any one of claims 1 to 27 which further comprises a viral component which is replaced with an equivalent component from a different serotype or is modified such that binding of said virus by antibodies pre-formed to the wild type component is reduced.

10

29. The modified virus of claim 28 wherein said viral component is a hexon protein.

15 30. A modified viral protein comprising a non-native polypeptide as defined in any one of claims 1 to 25.

31. A cell containing a modified virus or viral protein as defined in any of the claims 1 to 30.

20 32. A permissive cell for a modified virus as defined in any one of claims 1 to 29 which is capable of being cultured to propagate said modified virus.

25 33. A method for producing a modified virus as defined in any one of claims 1 to 29 in cell culture, comprising the steps of:

30 i) genetically modifying a virus to produce a modified virus comprising one or more non-native polypeptides, which polypeptide comprises one or more framework moieties each containing one or more binding moieties, which polypeptide is capable of being expressed in the cytoplasm and nucleus of a mammalian host cell in a conformation which is maintained in the absence of a ligand for said binding moieties, said conformation allowing said binding
35 moieties subsequently to bind with said ligand, and which

polypeptide is capable of transport though the nuclear membrane, wherein said modified virus has an altered tropism conferred by said binding moieties;

- 5 ii) infecting permissive cells with said modified virus;
iii) culturing said cells to produce the virus;
and
iv) harvesting, and optionally, purifying the modified virus produced.

10 34. A method of regulating the replication of a modified virus as defined in any one of claims 1 to 29 comprising the steps of:

- 15 i) constructing a modified virus such that a cleavage site is positioned between a binding moiety required for cell infection and the remainder of the recombinant viral component of which the binding moiety forms part; and,
ii) bringing said recombinant virus into contact with a cleavage agent or cleavage means capable of cleaving said binding moiety from said viral component and thereby
20 preventing the recombinant virus from undergoing further infection cycles.

25 35. A method of determining the suitability of a non-native polypeptide for use in the preparation of a viral vector by determining its solubility in a cell system.

36. A method as claimed in claim 35 which comprises the steps of:

- 30 i) expressing said non-native polypeptide or a viral component protein comprising said non-native polypeptide in permissive cells;
ii) subjecting the cells to lysis to produce a cell lysate;
iii) separating the soluble and insoluble fractions of the
35 cell lysate;

- - 155 - -

iv) analysing the soluble and insoluble fractions of the cell lysate for the content of said non-native polypeptide or viral component protein comprising said non-native polypeptide;

5 and,

v) comparing the relative content of said non-native polypeptide or viral component protein comprising said non-native polypeptide in the soluble and insoluble fractions

10

37. A method of assaying the solubility of a non-native polypeptide as defined in any one of claims 1 to 25 or a modified viral protein as defined in claim 30, which method comprises the steps of:

15 i) expressing said non-native polypeptide or a viral component protein comprising said non-native polypeptide in permissive cells;

ii) subjecting the cells to lysis to produce a cell lysate;

20 iii) separating the soluble and insoluble fractions of the cell lysate;

iv) analysing the soluble and insoluble fractions of the cell lysate for the content of said non-native polypeptide or viral component protein comprising said non-native polypeptide;

25 and,

v) comparing the relative content of said non-native polypeptide or viral component protein comprising said non-native polypeptide in the soluble and insoluble fractions.

30

38. The modified virus of any one of claims 1 to 29 for use in therapy.

35 39. Use of the modified virus of any one of claims 1 to

- - 156 - -

29 in the preparation of a medicament for the treatment of tumour cells or proliferating cells.

5 40. A pharmaceutical composition comprising a modified virus of any one of claims 1 to 29 and a pharmaceutically acceptable carrier or excipient.

Figure 1




Figure 2

A. WT Fiber



B. A7 EGF



C. A7 scFv C242



D. A7 scFv G250



E. A7 Z_{IgG}



F. A7 Affi IgA



G. A7 Z_{IgG}/Z_{IgA}



H. A7 Z_{IgG}/Z_{IgG}



I. A7 Z_{IgG} Xa Knob






Figure 3




FIGURE 4




FIGURE 5

BEST AVAILABLE COPY

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
31 January 2002 (31.01.2002)

PCT

(10) International Publication Number
WO 02/08263 A3

(51) International Patent Classification⁷: **C12N 15/86**,
15/861, 15/864, 15/867, A61K 35/76

(21) International Application Number: **PCT/GB01/03252**

(22) International Filing Date: **19 July 2001 (19.07.2001)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
GB 01 2014 19 July 2000 (19.07.2000) GB

(71) Applicant *for all designated States except US*: **GOT-A-GENE AB [SE/SE]**, Stena Center 1B, S-412 92 Göteborg (SE)

(71) Applicant *for GB only*: **GARDNER, Rebecca** [GB/GB] Frank B. Dehn & Co., 179 Queen Victoria Street, London EC4V 4EL (GB).

(72) Inventors; and

(75) Inventors/Applicants *(for US only)*: **LINDHOLM, Leif [SE/SE]**; Got-A-Gene AB, Stena Center 1B, S-412 92 Göteborg (SE). **NORD, Anna, Karin [SE/SE]**; Affibody AB, Björnåsvägen 21, S-113 47 Stockholm (SE). **BOULANGER, Pierre, Alain [FR/FR]**; Faculté de Médecine RTH Laennec, Laboratoire de Virologie et Pathogénèse Virale CNRS UMR 5537, 7, rue Guillaume Paradin, F-69372 Lyon Cedex 8 (FR).

(74) Agents: **GARDNER, Rebecca et al.**; Frank B. Dehn & Co., 179 Queen Victoria Street, London EC4V 4EL (GB).

(81) Designated States *(national)*: AE, AG, AL, AM, AT, AT (utility model), AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, CZ (utility model), DE, DE (utility model), DK, DK (utility model), DM, DZ, EC, EE, EE (utility model), ES, FI, FI (utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN.

[Continued on next page]

(54) Title: **MODIFIED VIRUS HAVING AN ALTERED TROPISM**

A. WT Fiber



B. A7 EGF



C. A7 scFv C242



D. A7 scFv G250



E. A7 Z_{IgG}



F. A7 Affi IgA



G. A7 Z_{IgG}/Z_{IgA}



H. A7 Z_{IgG}/Z_{IgG}



I. A7 Z_{IgG} Xa Knob



(57) Abstract: The present invention describes a modified virus comprising one or more non-native polypeptides, which polypeptide comprises one or more framework moieties each containing one or more binding moieties, which polypeptides is capable of being expressed in the cytoplasm and nucleus of a mammalian host cell in a conformation which is maintained in the absence of a ligand for said binding moieties, said conformation allowing said binding moieties subsequently to bind with said ligand, and which polypeptide is capable of transport through the nuclear membrane, wherein said modified virus has an altered tropism conferred by said binding moieties and the use of such viruses in the therapy, particularly in the treatment of tumours or other cancerous cells.

WO 02/08263 A3



MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

Published:

— with international search report

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(88) Date of publication of the international search report:

13 June 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

INTERNATIONAL SEARCH REPORT

national Application No

PCT/GB 01/03252

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/86 C12N15/861 C12N15/864 C12N15/867 A61K35/76

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, WPI Data, PAJ, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 20051 A (GENVEC INC ; KOVESDI IMRE (US); BROUGH DOUGLAS E (US); WICKHAM THOM) 5 June 1997 (1997-06-05)	1-15, 17, 19, 21, 22, 24-33, 35-40
Y	page 33, line 31 - page 34, line 17; examples 8, 19	16, 18, 20, 23, 34
X	WO 99 41359 A (UAB RESEARCH FOUNDATION) 19 August 1999 (1999-08-19)	1-15, 17, 19, 21, 22, 24-33, 35-40
Y	figure 1; examples 3, 4	16, 18, 20, 23, 34

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

6 March 2002

Date of mailing of the international search report

22/03/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Lanzrein, M

INTERNATIONAL SEARCH REPORT

national Application No

PCT/GB 01/03252

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 94 10323 A (EPENETOS AGAMEMNON ANTONIOU ;IMP CANCER RES TECH (GB); SPOONER ROB) 11 May 1994 (1994-05-11) page 6, line 10-25; figure 5; examples 1-4 ---	1-15,17, 25, 31-33, 38-40
X	WO 00 15823 A (GENVEC INC) 23 March 2000 (2000-03-23) page 5, line 34 -page 6, line 27; examples 5-10 ---	1-15,17, 25, 31-33, 38-40
Y	WO 95 19374 A (PHARMACIA AB ;NILSSON BJOERN (SE); NYGREN PER AAKE (SE); UHLEN MAT) 20 July 1995 (1995-07-20) page 6, line 3-32; claims 5,7,8 ---	16,18
Y	HOPPE JANS-JUERGEN ET AL: "A parallel three stranded alpha-helical bundle at the nucleation site of collagen triple-helix formation." FEBS (FEDERATION OF EUROPEAN BIOCHEMICAL SOCIETIES) LETTERS, vol. 344, no. 2-3, 1994, pages 191-195, XP001061304 ISSN: 0014-5793 abstract; figure 1 ---	20
Y	US 6 057 155 A (ROELVINK PETRUS W ET AL) 2 May 2000 (2000-05-02) column 18, line 8-20 ---	23,34
A	WO 00 12738 A (AVENTIS PHARMA SA ;LATTA MARTINE (FR); YEH PATRICE (FR); DEDIEU JE) 9 March 2000 (2000-03-09) examples 4-7 ---	1-40
A	WICKHAM T J: "Targeting adenovirus." GENE THERAPY., vol. 7, no. 2, January 2000 (2000-01), pages 110-114, XP001055392 ISSN: 0969-7128 page 112, left-hand column, paragraph 3 -page 113, right-hand column, paragraph 1 ---	1-40
P,X	WO 01 02431 A (LINDHOLM LEIF ;GOT A GENE (SE)) 11 January 2001 (2001-01-11) claims 12-16; figures 1,4 ---	1-40

-/--

INTERNATIONAL SEARCH REPORT

national Application No
PCT/GB 01/03252

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
T	<p>MAGNUSSON MARIA K ET AL: "Genetic retargeting of adenovirus: Novel strategy employing "deknobbing" of the fiber." JOURNAL OF VIROLOGY, vol. 75, no. 16, August 2001 (2001-08), pages 7280-7289, XP001056142 ISSN: 0022-538X the whole document -----</p>	1-40

INTERNATIONAL SEARCH REPORT

Information on patent family members

national Application No

PCT/GB 01/03252

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9720051	A	05-06-1997	US 5962311 A	05-10-1999
			US 5846782 A	08-12-1998
			US 5965541 A	12-10-1999
			AU 724189 B2	14-09-2000
			AU 1086897 A	19-06-1997
			AU 734515 B2	14-06-2001
			AU 4155597 A	06-03-1998
			BG 102554 A	30-04-1999
			BR 9612685 A	20-07-1999
			CZ 9801623 A3	14-10-1998
			EE 9800160 A	15-12-1998
			EP 0863987 A2	16-09-1998
			EP 0920524 A1	09-06-1999
			JP 2000516098 T	05-12-2000
			JP 11501219 T	02-02-1999
			NO 982417 A	06-07-1998
			PL 327038 A1	09-11-1998
			SK 72298 A3	02-12-1998
			WO 9720051 A2	05-06-1997
			WO 9807877 A1	26-02-1998
			AU 732770 B2	26-04-2001
			AU 4080497 A	06-03-1998
			EP 0920514 A1	09-06-1999
			HU 0001120 A2	28-07-2000
			JP 2001503250 T	13-03-2001
			WO 9807865 A1	26-02-1998
			US 6057155 A	02-05-2000
			US 6329190 B1	11-12-2001
WO 9941359	A	19-08-1999	AU 3294099 A	30-08-1999
			BR 9908018 A	24-10-2000
			CN 1297479 T	30-05-2001
			EP 1070118 A1	24-01-2001
			NO 20004563 A	13-09-2000
			WO 9941359 A1	19-08-1999
			US 6210946 B1	03-04-2001
WO 9410323	A	11-05-1994	AT 198769 T	15-02-2001
			DE 69329879 D1	22-02-2001
			DE 69329879 T2	07-06-2001
			DK 672158 T3	29-01-2001
			EP 1038967 A2	27-09-2000
			EP 0672158 A1	20-09-1995
			ES 2155074 T3	01-05-2001
			WO 9410323 A1	11-05-1994
			GB 2286593 A , B	23-08-1995
			JP 8506239 T	09-07-1996
			PT 672158 T	31-07-2001
			US 5885808 A	23-03-1999
WO 0015823	A	23-03-2000	AU 5819499 A	03-04-2000
			EP 1112372 A1	04-07-2001
			WO 0015823 A1	23-03-2000
WO 9519374	A	20-07-1995	AU 696186 B2	03-09-1998
			AU 1548795 A	01-08-1995
			CA 2181042 A1	20-07-1995
			EP 0739353 A1	30-10-1995

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 01/03252

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9519374	A		JP 9508016 T	19-08-1997
			NZ 278991 A	24-04-1997
			WO 9519374 A1	20-07-1995
			US 5831012 A	03-11-1998
<hr/>				
US 6057155	A	02-05-2000	US 5846732 A	08-12-1993
			US 5965541 A	12-10-1999
			US 6329190 B1	11-12-2001
			AU 724189 B2	14-09-2000
			AU 1086897 A	19-06-1997
			AU 732770 B2	26-04-2001
			AU 4080497 A	06-03-1993
			BG 102554 A	30-04-1999
			BR 9612685 A	20-07-1999
			CZ 9801623 A3	14-10-1993
			EE 9800160 A	15-12-1993
			EP 0863987 A2	16-09-1993
			EP 0920514 A1	09-06-1999
			HU 0001120 A2	28-07-2000
			JP 2001503250 T	13-03-2001
			JP 11501219 T	02-02-1999
			NO 982417 A	06-07-1993
			PL 327038 A1	09-11-1993
			SK 72293 A3	02-12-1993
			WO 9720051 A2	05-06-1997
			WO 9807865 A1	26-02-1993
<hr/>				
WO 0012738	A	09-03-2000	AU 5439399 A	21-03-2000
			CN 1314948 T	26-09-2001
			EP 1108047 A1	20-06-2001
			WO 0012738 A1	09-03-2000
<hr/>				
WO 0102431	A	11-01-2001	AU 6040100 A	22-01-2001
			WO 0102431 A1	11-01-2001

THIS PAGE BLANK (USPT